

Parity Doubling Among the Baryons

R. L. Jaffe, D. Pirjol, and A. Scardicchio

*Center for Theoretical Physics,
Laboratory for Nuclear Science and Department of Physics
Massachusetts Institute of Technology,
Cambridge, Massachusetts 02139*

Abstract. We study the evidence for and possible origins of parity doubling among the baryons. First we explore the experimental evidence, finding a significant signal for parity doubling in the non-strange baryons, but little evidence among strange baryons. Next we discuss potential explanations for this phenomenon. Possibilities include suppression of the violation of the flavor singlet axial symmetry ($U(1)_A$) of QCD, which is broken by the triangle anomaly and by quark masses. A conventional Wigner-Weyl realization of the $SU(2)_L \times SU(2)_R$ chiral symmetry would also result in parity doubling. However this requires the suppression of families of *chirally invariant* operators by some other dynamical mechanism. In this scenario the parity doubled states should decouple from pions. We discuss other explanations including connections to chiral invariant short distance physics motivated by large N_c arguments as suggested by Shifman and others, and intrinsic deformation of relatively rigid highly excited hadrons, leading to parity doubling on the leading Regge trajectory. Finally we review the spectroscopic consequences of chiral symmetry using a formalism introduced by Weinberg, and use it to describe two baryons of opposite parity.

INTRODUCTION

The possibility that excited hadrons occur in nearly degenerate pairs of opposite parity — a phenomenon known as parity doubling — has been considered from time to time since the 1960's. Early arguments were based on Regge theory [1]. There have been speculations that dynamical symmetries, not appearing in the underlying QCD Lagrangian, might be responsible for this phenomenon [2, 3]. Most recently it has been suggested that parity doubling can be explained by “restoration” of the underlying $SU(2)_L \times SU(2)_R$ chiral symmetry of QCD at high energies [4, 5, 6, 7, 8] and/or in specific sectors of the spectrum [9, 10].

In this paper we carry out a detailed analysis, both phenomenological and theoretical, of parity doubling among the baryons made of u , d , and s quarks. We do not consider mesons here because there is less data and more controversy. Given a concordance on meson resonances, our methods could easily be applied.

We address three questions: First, what is the experimental evidence for parity doubling? Second, could a symmetry of the underlying QCD Lagrangian be responsible for parity doubling? And third, even if the origins of parity doubling lie elsewhere, $SU(2)_L \times SU(2)_R$ is of fundamental importance for QCD, and is widely discussed in connection with spectroscopy. What, if any, are its implications for the classification of hadrons? On the first question, we find significant evidence for parity doubling among

non-strange baryons (nucleons and Δ s), but only weak evidence among hyperons (Σ s and Λ s) with strangeness (S) minus one. There is not enough data to carry out an analysis for Ξ s ($S = -2$) or Ω^- s ($S = -3$). Better information on the existence and classification of baryon resonances would help significantly here.

Turning to the possible origins of parity doubling, the most elegant explanations proposed make use of the fundamental symmetries of the QCD Lagrangian: chiral $SU(2)_L \times SU(2)_R$ and the singlet axial symmetry $U(1)_A$. Any underlying symmetry behind parity doubling must involve charges that transform as pseudoscalars, since the multiplets involve states of opposite parity. The natural candidates are the charges of the isospin axial currents, Q_5^a ($a = 1, 2, 3$), and the flavor singlet axial charge, Q_5 . $U(1)_A$ connects states of opposite parity, but the same flavor and spin. $SU(2)_L \times SU(2)_R$ charges change both parity and isospin, so multiplets in general contain states of opposite parity and different isospin.

Chiral symmetry restoration has been proposed as a mechanism for parity doubling for high excitations in the baryons spectrum [4, 5, 6, 7, 8]. This phenomenon was given the name “effective chiral restoration”, which we also use in the following for ease of comparison with the literature. It is equivalent to realizing chiral symmetry linearly on the excited states. As we pointed out in a recent paper [11], $SU(2)_L \times SU(2)_R$ chiral symmetry, when applied to the effective Lagrangian describing the interactions of excited baryons and matter, does not, by itself, lead to parity doubling. One can write down what appear to be Wigner-Weyl representations, where hadrons of equal mass and opposite parity transform into one another under the action of the $SU(2)$ axial charges. However $SU(2)_L \times SU(2)_R$ *invariant* operators that couple the would-be parity doublets to pions can be added to the effective Lagrangian describing these particles. These operators have the effect of removing the degeneracy of the previously degenerate states and spoiling any relationship between their axial charges. Non-linear field redefinitions make the situation clear (and do not change the physics): The only irreducible representations are multiplets of definite isospin and parity that transform according to the non-linear realization [12, 13, 14, 15]. To obtain degenerate pairs of opposite parity (and hence to achieve parity doubling) it is necessary to suppress the families of $SU(2)_L \times SU(2)_R$ *invariant* pion couplings that split them.

The existence of these operators and the need to suppress them dynamically has not been systematically examined in work on effective chiral symmetry restoration [4, 5, 6].

A somewhat different approach is taken in Refs. [9, 10], where hadrons are assigned to specific representations of $SU(2)_L \times SU(2)_R$ in the context of a linear sigma model. Chiral symmetry is subsequently broken when the sigma field takes a vacuum expectation value. By assigning hadrons to minimal (though not irreducible) $SU(2)_L \times SU(2)_R$ representations, and allowing only low dimension operators in the linear sigma model Lagrangian, a phenomenological framework for describing baryon resonances is constructed. In some examples, Refs. [9, 10] take parity doubling as an input when they choose representations and interactions. In fact the example we discuss at length in Sec. III was first introduced in Ref. [16] as a “mirror” model of opposite parity baryons. Because Refs. [9, 10] do not attempt either to marshal evidence for parity doubling in the data or explain it in terms of some more fundamental dynamics, we do not discuss them further here.

If “effective chiral symmetry restoration” is responsible for parity doubling, then

states, to the extent they are parity doubled, must decouple from pions. This prediction can be tested experimentally. Furthermore, only the simplest $SU(2)_L \times SU(2)_R$ multiplets involve only parity doubled pairs of states. Generic representations include states of different isospin as well as parity, which should be characteristic of this mechanism for parity doubling.

In contrast, the $U(1)_A$ symmetry of QCD is broken explicitly by a term in the quantum action induced by instantons [17] and, of course, by quark masses. Thus, its generator, Q_5 , the associated singlet axial charge, does not commute with the QCD Hamiltonian[17]. It is also not a symmetry of the QCD vacuum, since, as Coleman showed long ago[18], a charge that does not commute with the Hamiltonian cannot annihilate the vacuum. Thus $Q_5|0\rangle \neq 0$. Note, however, that this *does not* mean that $U(1)_A$ is “spontaneously broken”. In a theory like QCD with a non-perturbative $\langle \bar{q}q \rangle$ condensate in addition to instantons and quark masses, the question of whether a chiral symmetry is explicitly or spontaneously broken is *a matter of degree*. At one extreme, for zero up and down quark mass, $SU(2)_L \times SU(2)_R$ is clearly spontaneously broken. As an example of the other extreme, consider QED: Since $m_e \neq 0$, the axial $U(1)_A$ associated with the current $\bar{e}\gamma_\mu\gamma_5 e$ is not a symmetry of QED. Its charge does not commute with the Hamiltonian and does not annihilate the vacuum. In fact one can compute the $\bar{e}e$ condensate in perturbation theory. Specifically, in the $\overline{MS}$ scheme this is $\langle \bar{e}e \rangle_\mu = m^3/(4\pi^2) f(\alpha)$, where $f(\alpha) = 1 + \ell + \alpha/(2\pi)(5 + 5\ell + 3\ell^2) + O(\alpha^2)$, with $\ell = \log(\mu^2/m^2)$ and μ is the renormalization point [19]. Despite the fact that $\langle \bar{e}e \rangle \neq 0$, we say this symmetry is *explicitly broken*. In QCD with three colors, the importance of *explicit* $U(1)_A$ symmetry violation can be gauged by the magnitude of the η' mass, which comes entirely from explicit symmetry violation and is much greater than the scale of spontaneous symmetry breaking, f_π . Therefore, as discussed by 't Hooft[17], $U(1)_A$ behaves like an explicitly broken symmetry. This is one place where it is *not* useful to take the large N_c limit in QCD. There the situation becomes reversed because the source of explicit $U(1)_A$ symmetry violation disappears and the η' becomes a ninth pseudoscalar Goldstone boson (when $m_{u,d,s} = 0$). In that limit the $U(1)_A$ symmetry should be regarded as spontaneously broken. However, we live in a world with $N_c = 3$ and explicit breaking of $U(1)_A$ dominates.

Like any *explicitly broken* symmetry, if the matrix elements *that violate* $U(1)_A$ symmetry are small in a sector of the spectrum, some of its consequences will follow. In the case at hand, parity doubling will be seen in a sector of the spectrum if the matrix elements of the divergence of the flavor singlet axial current are dynamically suppressed in that sector [20]. Other consequences of the symmetry are not obtained: for example the parity doubled states do not necessarily form representations of $U(1)_A$. For the sake of clarity, we refer to this phenomenon as “dynamical suppression of $U(1)_A$ symmetry violation” rather than “ $U(1)_A$ symmetry restoration”.¹

To summarize, either chiral symmetry or $U(1)_A$ could be responsible for parity doubling. In both cases extra dynamical mechanisms must be at work. For $SU(2)_L \times SU(2)_R$, *chirally invariant* operators must be suppressed. If so, hadrons form chiral multiplets,

¹ In an earlier version of this paper we argued that suppression of $U(1)_A$ violation would be less likely among strange baryons. We no longer believe this to be required. We thank A. Manohar for discussions on this point.

not necessarily limited to parity doublets, and they decouple from pions. $U(1)_A$ is more conventional: parity doubling can result if operators that *violate the symmetry* are suppressed.

We also briefly explore some other proposed dynamical origins for parity doubling. We consider the possibility that parity doubling is not directly associated with a symmetry of the underlying Lagrangian, but instead is a consequence of an intrinsic deformation of excited hadrons. If baryons can be accurately described as intrinsically deformed systems, if the intrinsic state spontaneously violates parity, and if the intrinsic state is rigid, *i.e.* the matrix element between the state and its parity image is small, then when collectively quantized, the spectrum would exhibit parity doubling. This is, in fact, an old idea [1, 2, 22], developed in early geometrical models of baryons. The arguments in support of this mechanism seem to apply best to states of high angular momentum, and there is evidence for parity doubling at high J among non-strange baryons. However we also find strong evidence for parity doubling among baryons with $J = 1/2$ and $3/2$, which is hard to motivate from this point of view. We also discuss the arguments given by Shifman [8] that the chiral invariance of perturbative QCD, valid at short distances, together with certain dynamical assumptions motivated by the $N_c \rightarrow \infty$ (N_c is the number of colors in QCD) limit, lead to parity doubling among mesons, and perhaps by extension, baryons.

Finally, although $SU(2)_L \times SU(2)_R$ is in general not apparent in the spectrum, it is not devoid of implications for the spectrum of baryons. Weinberg has shown that hadrons can be classified into multiplets under an $SU(2)_L \times SU(2)_R$ symmetry where the axial generators are associated with pion-hadron vertices, provided hadron-hadron scattering amplitudes obey certain constraints at asymptotic energies [23, 24]. Weinberg’s “mended” chiral symmetry does not naturally lead to parity doubling, but it can accommodate it for special choices of the hadrons’ assignments into $SU(2)_L \times SU(2)_R$ representations. We explore the possible spectroscopic consequences of chiral symmetry in the context of Weinberg’s work on “mended symmetries”, and comment on the relation of this approach to the contracted spin-flavor symmetry emerging in the large N_c limit [25, 26].

The remainder of the paper is organized as follows. In the next section we examine the experimental situation. Despite all the theoretical work, we know of no systematic attempt to quantify the evidence for parity doubling. Typically papers on the subject include graphs or tables where the reader is invited to see some evidence for a correlation between states of opposite parity (for a recent example, see Ref. [7]). It is difficult, however, to factor in the (often large) widths of states and the fact that some states are much better established than others. Also it is not clear which state to pair with which. Some previous attempts to quantify the evidence for (or against) parity doubling can be found in Refs. [27, 28].

We introduce a measure of the correlation between states of opposite parity, $\Omega(I, J, S)$, that has some statistical basis. Specifically, we test the hypothesis that two functions, positive and negative parity spectral densities for each isospin (I), strangeness (S), and spin (J), are identical within a tolerance σ , which measures the extent of symmetry breaking. We construct the spectral functions for each I and J from the baryons accepted by the Particle Data Group [29], including widths, and weighting resonances to reflect their reliability. We have no *a priori* standard with which to compare Nature’s value of

$\Omega(I, J, S)$. Instead we construct a set of “alternative realities” by scrambling the parities of the known states. As an ensemble, this set is free from correlations, so the comparison between $\Omega(I, J, S)$ computed over this ensemble with the value obtained from Nature’s assignment of parities gives a measure of the significance of the correlation. The method and our results are discussed in detail in Section II.

In Section III we look at the fundamental symmetries of QCD to see which, if any, could be responsible for the phenomenon of parity doubling. We review the realizations of chiral symmetry and briefly summarize the arguments of our recent paper [11] on the way that massless Goldstone bosons transforming non-linearly undermine the predictions of linearly realized chiral symmetry for baryon masses and couplings. We go on to state the conditions on the chiral effective Lagrangian that must be satisfied in order to obtain parity doubling, irrespective of the underlying dynamical mechanism responsible for it. We describe the phenomenon of “dynamical suppression of $U(1)_A$ breaking”, and show that it can lead to parity doubling in a sector of the Hilbert space of QCD. We also summarize and review arguments for parity doubling from chiral symmetry restoration at short distances, and due to intrinsic hadron deformation. For completeness we summarize the spectroscopic predictions of Weinberg’s approach in Section IV, where we introduce a new example that reproduces naturally some of the predictions of $SU(2)_L \times SU(2)_R$ restoration, without fine tuning of the coefficients in the chiral Lagrangian. Finally in Section V we summarize our results and discuss areas for further work.

WHAT IS THE EXPERIMENTAL EVIDENCE?

In this section we evaluate the experimental evidence for parity doubling among the baryons. We have chosen to study the baryons as listed by the Particle Data Group[29]. There is still much uncertainty concerning the spectrum of light (u , d , s) baryons and mesons, and other authors have chosen to study different data[4, 5]. We chose the PDG listings because they provide a critical summary of many experiments over many decades, and also because they treat baryon states uniformly, assigning masses, widths, and “reliabilities” using their famous “star” system. Data on the meson spectrum accepted by the PDG is less complete, more controversial and lacks the reliability assignments. This point requires some discussion. The PDG has assigned reliabilities to 44 non-strange baryon resonances[30]. For each angular momentum there are *two* isospin channels, $I = 1/2$ (nucleon) and $I = 3/2$ (Δ). The PDG non-strange meson listings are harder to interpret. The “Meson Summary Table” lists (coincidentally) 44 well-established non-strange mesons. However those states are distributed over *four* channels, $I = 0$ and $I = 1$ with $C = \pm 1$. Many more states appear in the full meson listings: 70 states can be found in the non-strange meson listings and another 96(!) can be found under the heading “Other Light Unflavored Mesons”. The latter list includes some that have been seen in only one experiment and some that other groups have claimed to exclude. Recently, Bugg[31] has made an independent analysis of the meson spectrum including an attempt at assigning reliabilities, with results quite different from the PDG analysis. It would be very interesting to repeat our statistical analysis on a well-defined

set of meson resonances.

We find that there is reasonably strong evidence for parity doubling among the non-strange baryons ($I = 1/2$ nucleons and $I = 3/2$ Δ s), and weaker, inconclusive evidence among the $S = -1$ baryons ($I = 0$ Λ s and $I = 1$ Σ s). There is not enough data on cascades ($S = -2$) or Ω s ($S = -3$) to perform our analysis. The results are summarized in Figs. 2, 3, 4 and 5, which should be consulted after an explanation of our method. It will be clear that the study of parity doubling is hampered by our incomplete knowledge of baryon resonances, especially those with strangeness. Certainly a new experimental attack on hadron spectroscopy could greatly improve matters.

Despite many papers on the subject over many years, we know of no attempt to make a (semi-) quantitative evaluation of the evidence for parity doubling. Typically authors simply show the spectrum of states with the same flavor and spin quantum numbers and opposite parity, and rely on the eye of the reader to recognize a propensity for states of opposite parity to cluster nearby in mass. Such an “eyeball” approach has several shortcomings,

- Baryon resonances have large widths that are hard to display graphically.
- Different resonances have different statistical significance. Overlap of two well established resonances should be given more weight than overlap of two doubtful states.
- There is no reason to expect parity doubling to be an exact symmetry, so spectra must be compared with an allowance for symmetry breaking.
- Because of symmetry breaking and resonance widths, in channels with many states it is often not clear which states are to be compared.

Recently Glazman and Klemp[27, 28] have made more quantitative analyses, which however still suffer from some of the shortcomings listed above. In our analysis we will address all of these difficulties.

We know of no way to obtain a rigorous measure of the significance of parity doubling in the physical spectrum, which perhaps explains the absence of work in this direction. However we do believe it is possible to improve on the eyeball. To do this one needs two ingredients: 1) A measure of how much correlation there is between negative and positive parity states; and 2) A control set of “alternative worlds” on which the correlation can be measured and compared with our own.

A realistic approach to generating control spectra has to respect several features of the baryon resonance spectrum: a) states are sparse at low mass, where there is no evidence for parity doubling; b) the data in each channel show roughly equal numbers of positive and negative parity states; c) reliabilities decrease and widths increase in general with resonance mass. Reference spectra that ignore these features would not give fair comparisons with the real world. We do not know an algorithm that would generate *a priori* satisfactory comparison spectra free of these problems.

Instead we have chosen to use the physical spectrum itself to generate comparisons. Specifically, in each channel of definite J , I , and S , we keep the masses, widths, and reliabilities fixed, and reshuffle the the parities, keeping the number of positive and negative parity states fixed. This method generates comparison spectra which respect the characteristics listed in the previous paragraph, but which, as an ensemble, is free

from any additional parity correlation beyond the desirable feature that the number of states of opposite parities is roughly equal in each channel. Comparing the real world with these alternative realities tests whether the correlation between states of opposite parity in the real world beats a random ensemble with otherwise similar spectroscopic features.

We consider “channels” of specific total angular momentum, J , isospin, I , and strangeness, S . In each channel we define positive and negative parity spectral functions, $\rho_{IJS}^{\pm}(m)$, which we take to be a sum over normalized Breit-Wigner resonances with masses and widths taken from the PDG². Specifically, we include only the states listed in the “Note on Nucleon and Δ Resonances”[30]. We weigh the resonances by a confidence factor, W_j , proportional to the number of stars designated by the PDG, because we believe more weight should be given to correlations between well established states,

$$\begin{aligned}\rho_{IJS}^{+}(m, \{C\}) &= \sum_j \frac{W_j}{2\pi} \frac{\Gamma_j}{(m - m_j)^2 + \Gamma_j^2/4} C_j \\ \rho_{IJS}^{-}(m, \{C\}) &= \sum_j \frac{W_j}{2\pi} \frac{\Gamma_j}{(m - m_j)^2 + \Gamma_j^2/4} (1 - C_j)\end{aligned}\quad (1)$$

The sum ranges over *all* states with a definite J , I , and S , and $W_j = 1.0, 0.75, 0.50, 0.25$ for 4^* , 3^* , 2^* , and 1^* resonances respectively. We have also performed the analysis with an exponential weighting, $W_j = 1.0, 0.50, 0.25, 0.125$ respectively. The effect on our results is negligible except for the Σ s ($I = 1, S = -1$), where the shape of the distribution shifts significantly enough to warrant comment (see discussion below). The factor C_j assigns a parity to the j^{th} state: $C_j = +1$ for positive parity, and $C_j = 0$ for negative parity. Of course, Nature prescribes a parity for each state as a set of C_j ’s. Notice that the $\rho_{IJS}^{\pm}$ are normalized,

$$\int_{-\infty}^{\infty} dm \rho_{IJS}^{\pm}(m, \{C\}) \equiv N_{IJS}^{\pm}(\{C\}) = \sum_j W_j \begin{cases} C_j & \text{for } \rho^{+} \\ (1 - C_j) & \text{for } \rho^{-} \end{cases} \quad (2)$$

so the norm measures the number of states weighted by our confidence that they are real.

We do not expect parity doubling to be an exact symmetry of QCD. Instead we expect it to be approximately valid, parameterized by some symmetry breaking scale σ , which is expected to be of order (say) 50–200 MeV. Our aim, then, is to test the hypothesis that the masses of positive and negative parity states are compatible within fluctuations with a variance σ . A simple indicator we can use to test the above hypothesis is

$$\Omega_{IJS}(\{C\}) = \int dm_1 dm_2 \rho_{IJS}^{+}(m_1, \{C\}) \operatorname{erfc}\left(\frac{|m_1 - m_2|}{\sigma}\right) \rho_{IJS}^{-}(m_2, \{C\}) \quad (3)$$

where $\operatorname{erfc}(z) = \frac{2}{\sqrt{\pi}} \int_z^{\infty} dt e^{-t^2}$ is the complementary error function. $\operatorname{erfc}(0) = 1$ and $\operatorname{erfc}(z) \rightarrow 0$ when $z \rightarrow \infty$. Eq. (3) then can be interpreted as follows. For a given m_1 and

² We do not include the quoted uncertainties in masses and widths since they are generally smaller than the widths of the states, and including them would complicate our analysis without adding much value.

m_2 , $\text{erfc}(|m_1 - m_2|/\sigma)$ is the probability that a gaussian random number μ with average m_1 and variance σ gives a $|\mu - m_1| \geq |m_2 - m_1|$, i.e. the probability that m_2 is closer to m_1 than a randomly generated number (with variance σ). We can interpret σ as the order of magnitude of the breaking of the symmetry (the argument is symmetric in $m_2 \leftrightarrow m_1$). The possible values of m_1 and m_2 are then distributed according to $\rho_{IJS}^\pm(m_{1,2}, \{C\})$, normalized not to unity, but to $N_{IJS}^\pm(\{C\})$. $\Omega_{IJS}(\{C\})$ measures the correlation of the spectral functions with a “forgiveness factor” σ . Ω is going to be larger in channels with more states, greater reliability, and, of course, better correlation between states of opposite parity. Since $\text{erfc}(x) \leq 1$ the maximum possible value of Ω_{IJS} occurs if the typical separations $|m_i - m_j|$ and widths Γ_i are much smaller than the chosen σ . In that case we can set $\text{erfc}(|m_1 - m_2|/\sigma) \approx 1$ for all m_1 and m_2 and the two integrals decouple giving and $\Omega_{IJS}^* \equiv \text{Max}[\Omega_{IJS}(\{C\})] = N_{IJS}^+(\{C\})N_{IJS}^-(\{C\})$. Some other features of Ω_{IJS} are worth noting: There is no penalty for the absence of parity doubling low in the spectrum. A state like the nucleon with no parity partner simply does not contribute. Any overlapping states of opposite parity (and the same IJS) contribute to Ω_{IJS} . No choice of multiplet assignments is needed. The contribution of broad states is insensitive to shifts correlations on scales small compared to their widths.

We do not know how to judge the *a priori* significance of the experimental value of Ω_{IJS} because the underlying distributions $\rho_{IJS}^\pm$ are not normalized to unity, but instead weighted by the number and reliability in that channel — features that we regard to be important in the analysis. So we are led to check the value of Ω realized in Nature against an ensemble of control spectra generated by scrambling the parities among the otherwise fixed states for each J, I , and S . The crucial feature of these control spectra is that the parities are reassigned randomly. So the comparison between Nature and the control set frames the question whether the parities of the states with a given I, J, S are more correlated than a random assignment of parities.




FIGURE 1. The value of Ω for Nucleons ($I = 1/2, S = 0$) with $J = 1/2$ for varying σ . The value of Ω with the Nature’s parity assignments is the red line.

Fig. 1 shows the results in a typical channel, nucleons with $J = 1/2$. $\Omega_{1/2,1/2,0}$ is plotted for Nature’s parity assignments (in red) and the control sample (in gray), for a range of σ . The ordering of instances turns out not to depend significantly on σ , so we

lose little by choosing a fixed value of σ for the rest of the analysis. We have chosen $\sigma = 125$ MeV, though we cannot attribute any physical significance to this choice. Fig. 2 presents the data for nucleons and Δ s (for $J \leq 7/2$) and Fig. 3 shows the data for Σ s and Λ s.




FIGURE 2. The value of Ω at $\sigma = 125$ MeV for nucleons (left) and Δ 's (right) of different spins J . The value of Ω with Nature's parity assignments is marked with a red circle.




FIGURE 3. The value of Ω at $\sigma = 125$ MeV for Σ 's (left) and Λ 's (right) of different spins J . The value of Ω with Nature's parity assignments is marked with a red circle.




FIGURE 4. Histograms of frequency of the value of Ω in all the possible realizations of parity assignments in the nucleons (left) and Δ 's (right). All spins less than or equal $\frac{7}{2}$ are included. The value of Ω with the parities as observed in Nature is indicated with a star. The area on the left of the bar represents the 'confidence level' (C.L.) attached to the given Ω . For nucleons we have 95% C.L. and for the Δ 's 86% C.L.




FIGURE 5. Histograms of frequency of the value of Ω in all the possible realizations of parity assignments in the Σ (left) and Λ (right). All spins $J = 1/2, \dots, 7/2$ are included. The value of Ω with the parities as observed in Nature is indicated with a star. The area on the left of the bar represents the ‘confidence level’ (C.L.) attached to the given Ω . For Σ we have 60% C.L. and for the Λ 65% C.L. .

Data are most abundant for nucleons and Δ s, and for these two channels Nature shows typically more correlation than the alternative realities. To give a quantitative answer, in terms of confidence level, we construct a histogram of the frequency of a given value of Ω . Because the Ω_{IJS} is normalized in a way that reflects the significance of each channel, it makes sense to combine data for sets of channels, thereby obtaining a larger control ensemble. This has been done in Fig. 4 for nucleons and for Δ s. From these figures it is clear that although a small number of parity reassignments would do better, Nature’s correlation between states of opposite parity is quite striking. In numbers the ‘confidence level’ (the area on the left of the starred bin in Figure 4) for the value of Ω in the Nature is 95% for the nucleons and 86% for the Δ s.

The evidence for parity doubling among Λ s and Σ s, channel by channel, is shown in Fig. 3, and the statistic is summed over all $S = -1$ channels in Fig. 5. For the strange baryons we find much smaller values of the confidence level for Ω : 58% for the Σ s and 72% for the Λ s. We conclude that it is not possible to make a definitive statement in this sector. There may be some correlation, but more data are needed. Fig. 6 shows histogram using the “exponential” weighting factor for the Σ s. Here, the change in weighting makes a difference: the signal for parity doubling is stronger when well established resonances are more strongly weighted.

POSSIBLE ORIGINS OF PARITY DOUBLING

In this section we examine some possible explanations for the phenomenon of parity doubling among light quark baryons. In the classical limit, the Lagrangian of QCD with n_f light quark flavors has the symmetry

$$SU(n_f)_L \times SU(n_f)_R \times U(1)_A \times U(1)_V \quad (4)$$




FIGURE 6. Frequency histogram of the value of Ω for the Σ 's using the “exponential” weighting of the significance of states. Compare with Fig. 5, left box.

This consists of the product of chiral symmetry, $SU(n_f)_L \times SU(n_f)_R$, with the singlet axial symmetry, $U(1)_A$, and the baryon number invariance $U(1)_V$. Our understanding of QCD is profoundly shaped by these symmetries, so in the following we review their possible role in connection with the phenomenon of parity doubling. Since we are studying light quark spectroscopy but wish to keep track of strange quark mass effects, we take $n_f = 2$.

We start by reviewing the implications of chiral symmetry for the masses and couplings of hadrons. This part also introduces the notation to be used in the remainder of the paper. We also briefly review the arguments we presented in Ref. [11] that parity doubling does not follow from the underlying $SU(2)_L \times SU(2)_R$ symmetry of QCD. That symmetry can only be restored in the spectrum if classes of $SU(2)_L \times SU(2)_R$ invariant operators are suppressed for some dynamical reason.

Next we consider the possibility that parity doubling is a consequence of dynamical suppression of flavor singlet axial symmetry violation. As is usually the case with explicitly broken symmetries, the symmetry can be approximately realized in the spectrum if the matrix elements of operators that violate the symmetry are suppressed.

We then discuss two other possibilities: first Shifman's attempt to relate parity doubling to chiral symmetry restoration at short distances in QCD, and second the possibility that parity doubling is associated with an intrinsic deformed baryon state, a mechanism familiar from nuclear and molecular physics.

Effective restoration of $SU(2)_L \times SU(2)_R$

The Lagrangian of QCD with n_f massless quarks has an exact chiral $SU(n_f)_L \times SU(n_f)_R$ symmetry, with generators given by the vector T^a and axial X^a charges. The generators satisfy the commutation relations

$$[T^a, T^b] = i f_{abc} T^c, \quad [T^a, X^b] = i f_{abc} X^c, \quad [X^a, X^b] = i f_{abc} T^c, \quad (5)$$

where $a, b, c = 1, \dots, n_f^2 - 1$. For the case of $n_f = 2$ of most interest here, $f_{abc} = \epsilon_{abc}$. The vector and axial isospin currents are exactly conserved, $\partial^\mu j_\mu^a = \partial^\mu j_{5\mu}^a = 0$. The

generators have different transformation properties under parity: the vector generators, T^a , commute with parity, but the axial charges anticommute $\{X^a, P\} = 0$. This suggests that parity is embedded nontrivially in representations of the chiral group, in a way which could reproduce the observed parity doubling in the hadron spectrum.

It is instructive at the outset to review the consequences of current conservation for the spectrum. Conservation of the vector current requires hadrons to lie in degenerate multiplets of definite isospin. The consequences of axial current conservation are embodied in generalized Goldberger-Treiman relations. Consider two hadrons of identical flavor and spin but opposite parity — without loss of generality we can consider baryons of spin and isospin $1/2$ — and expand the matrix elements $j_{5\mu}^a$ in terms of form factors,

$$\langle B_+ | j_{5\mu}^a | B_- \rangle = \bar{u}(p, s) (\gamma_\mu g_A(q^2) + q_\mu g_P(q^2) + i\sigma_{\mu\nu} q^\nu g_M(q^2)) t^a u(p', s') \quad (6)$$

where $q^\mu = p^\mu - p'^\mu$. Conservation of the axial current demands that $q^\mu \langle B_+ | j_{5\mu}^a | B_- \rangle = 0$, which gives

$$\Delta m g_A(\Delta m^2) + \Delta m^2 g_P(\Delta m^2) = 0 \quad (7)$$

for states at rest ($\vec{q} = 0$), where $\Delta m = m_+ - m_-$. $g_P(q^2)$ has a pion pole at $q^2 = 0$ with residue $f_\pi g_{\pi B_+ B_-}$, where $g_{\pi B_+ B_-}$ is the s -wave $B_+ B_- \pi$ coupling constant. Then for small Δm ,

$$\Delta m g_A + f_\pi g_{\pi B_+ B_-} + O(\Delta m^2) = 0 \quad (8)$$

where $g_A = g_A(0)$.

Suppose for some reason that B_+ and B_- were degenerate. Then $\Delta m = 0$ and eq. (8) requires $g_{\pi B_+ B_-} = 0$ — the pion must decouple at zero momentum. The argument goes through for states of different isospin with $\Delta I = 1$, as well as for mesons and strange and heavy hadrons. Thus any states of the same spin, isospin, *etc.*, but opposite parity that have the same mass must decouple from pions at zero momentum. This is a consequence of axial current conservation. So chiral symmetry does not guarantee the states to be degenerate, but if they are, it forces them to decouple from pions at zero momentum, *i.e.* in the s -wave.

The thesis of Refs. [9, 10, 4, 5, 6] is that hadrons show parity doubling because at masses of order $1.5 - 3$ GeV (where the data show evidence of parity doubling) chiral symmetry is approximately restored in the hadron spectrum. By this they mean that the symmetry is realized (approximately) in Wigner-Weyl mode with the appearance of (approximately) degenerate multiplets that transform into one another linearly under chiral rotations.

The irreducible representations of $SU(2)_L \times SU(2)_R$ are labelled by the Casimirs of the left/right-handed group factors (I_L, I_R) which are related to the usual isospin and its axial partner by $I_{L,R}^a = T^a \mp X^a$. The representation (I_L, I_R) contains $(2I_L + 1)(2I_R + 1)$ states, with total isospin taking all possible values compatible with $|I_L - I_R| \leq I \leq I_L + I_R$. Parity exchanges the left and right-handed isospins $I_L \leftrightarrow I_R$, such that these irreducible representations are in general not parity invariant, unless $I_L = I_R$. We are interested in the baryon states such as N, Δ with isospins $1/2$ and $3/2$. These states can be contained only in chiral multiplets, $(1/2, 0)$, $(1/2, 1)$, $(3/2, 0)$, $(3/2, 1)$, $\dots$, which are not parity invariant. It is easy to form parity eigenstates by taking sums and differences of mirror

chiral representations

$$H_+ \sim (I_L, I_R) \oplus (I_R, I_L), \quad H_- \sim (I_L, I_R) \ominus (I_R, I_L) \quad (9)$$

In this example, $H_\pm$ represents parity-even(odd) hadrons. Therefore a Wigner-Weyl representation of $SU(2)_L \times SU(2)_R$ generically includes degenerate multiplets of both positive and negative parity and a range of isospins.

In addition to degeneracy, the linear realization of the $SU(2)_L \times SU(2)_R$ predicts relations between axial and vector charges typical of a non-Abelian symmetry. In Ref. [11] we showed that decoupling of pions implied by eq. (7) in the parity doubling limit is not a curiosity. Instead hadrons can only form Wigner-Weyl representations of $SU(2)_L \times SU(2)_R$ if they *completely decouple from pions*. Indeed this is just the mechanism proposed in Refs. [4, 5, 6], where it is argued that states high in the spectrum are generically insensitive to the effects of spontaneous symmetry breaking. Whether or not one accepts these arguments, this dynamical picture makes many predictions that can be tested. First and foremost is the prediction that the parity doubled states should decouple from pions, not only in the s -wave at zero momentum as required by the Goldberger-Treiman relation, eq. (7), but in higher partial waves as well. Also, the arguments of Ref. [4, 5, 6] in the parity doubled domain of the spectrum apply whether or not the particular states are degenerate. All the pion transition matrix elements should be small to the extent that chiral symmetry is “restored”. Second, larger multiplets including states of different isospin should be found in the spectrum. Glozman and Klempt[27, 28] have made some initial attempts to classify states into such multiplets, with varying amounts of success. Third, strange and heavy hadrons should show the same phenomenon — the emergence of explicit $SU(2)_L \times SU(2)_R$ symmetry high in the spectrum — since the heavy quarks are merely spectators to this dynamical mechanism. Finally, the phenomenon should be generic. If some well established states high in the spectrum *cannot* be classified into representations of $SU(2)_L \times SU(2)_R$, it is evidence that other dynamics is at work.

For the sake of completeness, we review the argument of Ref. [11] that pions must decouple from states that exhibit the full structure of $SU(2)_L \times SU(2)_R$ symmetry. Parity invariance of QCD requires that physical states are parity eigenstates. As explained above, this can be realized only if the physical states transform according to sums and differences of chiral representations as in eq. (9). Thus the combined consequence of chiral symmetry realized in WW mode and parity invariance of QCD, is the appearance of parity doubling in the hadron spectrum.

However, we know that in Nature chiral symmetry is realized in Nambu-Goldstone mode with the appearance of Goldstone bosons. Actually two distinct possible NG realizations have been discussed in the literature, distinguished by the invariance of the vacuum. In the usual realization $SU(2)_L \times SU(2)_R$ breaks to isospin. Long ago, Dashen pointed out that $SU(n_f)_L \times SU(n_f)_R$ could break to $SU(n_f)_V \times Z_{n_f}^A$, where $Z_{n_f}^A$ is the center of axial $SU(n_f)$ [32]. We consider the usual scenario first, and return to Dashen’s alternative at the end of this subsection.

Consider for definiteness two spin-1/2 baryons of positive (B_+) and negative (B_-)

parity, and unspecified isospin I^3 . Assume that they transform linearly under chiral symmetry

$$[T^a, B_+^i] = -t_{ij}^a B_+^j, \quad [T^a, B_-^i] = -t_{ij}^a B_-^j \quad (10)$$

$$[X^a, B_+^i] = t_{ij}^a B_-^j, \quad [X^a, B_-^i] = t_{ij}^a B_+^j \quad (11)$$

with i, j isospin indices. This corresponds to the $SU(2)_L \times SU(2)_R$ representations $B_+ \sim (I, 0) + (0, I)$ and $B_- \sim (I, 0) - (0, I)$. The most general chirally and parity invariant effective Lagrangian *containing only* $B_\pm$, reads

$$\mathcal{L} = \bar{B}_+ i \not{\partial} B_+ + \bar{B}_- i \not{\partial} B_- - m_0 (\bar{B}_+ B_+ + \bar{B}_- B_-) + \dots \quad (12)$$

where the ellipses denote terms of higher order in the chiral expansion. The two hadrons are degenerate, in agreement with the arguments of [6]. In addition, the matrix elements of the axial current are also fixed by the symmetry, via the Noether theorem.

So far we have neglected the possible coupling of B_+ and B_- to pions. The pion field transforms nontrivially under the chiral group. This has the consequence that additional operators, constructed from the baryon fields $B_\pm$ and π^a , can be added to the effective Lagrangian, eq. (12). For example, using Weinberg's choice[13, 14] for the pion non-linear transformation under axial rotations,

$$[X^a, \pi^b] = -i f^{ab}(\pi) = -i \left(\delta^{ab} \frac{1}{2} (1 - \vec{\pi}^2) + \pi^a \pi^b \right), \quad (13)$$

(where we expressed the pion field in units of $f_\pi = 131$ MeV) another dimension three, chirally invariant operator can be added to eq. (12),

$$\delta \mathcal{L} = m_1 \left(\bar{B} \frac{-1 - \pi^2}{1 + \pi^2} B - \bar{B} \frac{-4i\pi^a t^a}{1 + \pi^2} B' - (B \leftrightarrow B') \right)$$

Rediagonalizing the terms quadratic in $B_\pm$ we see that the new term has the effect of splitting the degeneracy: the new mass eigenvalues are $m_0 \pm m_1$. m_1 is the coupling constant set to zero by the generalized Goldberger-Treiman relation if the hadrons B_+ and B_- are to be degenerate.

The effects of pion couplings are most conveniently displayed by redefining the baryon fields,

$$\tilde{B}_+ = \frac{B_+ - 2i\pi^a t^a B_-}{\sqrt{1 + \pi^2}}, \quad \tilde{B}_- = \frac{B_- - 2i\pi^a t^a B_+}{\sqrt{1 + \pi^2}} \quad (14)$$

The new fields have also positive ($\tilde{B}_+$) and negative ($\tilde{B}_-$) parity, but transform nonlinearly under chiral rotations. Specifically,

$$[X^a, \tilde{B}_{\pm i}] = v_0(\pi^2) \epsilon_{abc} \pi^c t_{ij}^b \tilde{B}_{\pm j}, \quad (15)$$

³ The example which follows is equivalent to the “mirror model” presented in Ref. [16] in the context of the linear sigma model.

with $v_0(\pi^2)$ a function which depends on the particular definition of the pion field. We use here and in the following the definition adopted in Ref. [14], which corresponds to $v_0(x) = 1$. Most importantly, the new states $\tilde{B}_\pm$ do not transform into one another under chiral transformations. Instead of taking $B_+ \leftrightarrow B_-$ as in eqs. (10), the axial chiral rotation of the fields defined in eq. (14) transforms each of $\tilde{B}_i$ into itself, plus a number of pions. This agrees with the intuitive idea that an axial rotation does not transform hadrons within a chiral multiplet, but just rotates their isospins and creates pions from vacuum.

It is easy to construct the most general chirally invariant effective Lagrangian describing the dynamics of the baryons $\tilde{B}_\pm$ following the usual rules of chiral effective field theory [12, 13, 14]. When rewritten in terms of the original fields, $B_\pm$, we discover several other possible chirally invariant terms, allowed when pions transform nonlinearly, but omitted from eq. (12). They take the simplest form when expressed in terms of the $\tilde{B}_\pm$ fields, and are given by

$$\begin{aligned} \mathcal{L} = & \bar{\tilde{B}}_+(i\cancel{\partial} - \epsilon^{abc}\pi^a\cancel{D}\pi^bt^c)\tilde{B}_+ - \bar{\tilde{B}}_+(\cancel{D}\pi^a)t^a\tilde{B}_- \\ & + (\tilde{B}_+ \leftrightarrow \tilde{B}_-) - (m_0 - m_1)\bar{\tilde{B}}_+\tilde{B}_+ - (m_0 + m_1)\bar{\tilde{B}}_-\tilde{B}_- + \delta\mathcal{L}_2 \end{aligned} \quad (16)$$

where $D_\mu\pi^a = 2\partial_\mu\pi^a/(1+\pi^2)$ is the covariant derivative of the pion field. The additional terms $\delta\mathcal{L}_2$ contain three operators which couple the pions derivatively to the baryons

$$\begin{aligned} \delta\mathcal{L}_2 = & c_2[\bar{\tilde{B}}_+(\cancel{D}\pi^a)\gamma_5 t^a\tilde{B}_+ + \bar{\tilde{B}}_-(\cancel{D}\pi^a)\gamma_5 t^a\tilde{B}_-] \\ & + c_3[\bar{\tilde{B}}_+(\cancel{D}\pi^a)t^a\tilde{B}_- + \bar{\tilde{B}}_-(\cancel{D}\pi^a)t^a\tilde{B}_+] \\ & + c_4[\bar{\tilde{B}}_+(\cancel{D}\pi^a)\gamma_5 t^a\tilde{B}_+ - \bar{\tilde{B}}_-(\cancel{D}\pi^a)\gamma_5 t^a\tilde{B}_-] \end{aligned} \quad (17)$$

Note that the baryons $\tilde{B}_\pm$ are not degenerate, but are split by $2m_1$. Similar results are obtained for the matrix elements of the axial current, which is obtained from the Noether theorem

$$\begin{aligned} A_\mu^a = & (c_2 + c_4)\bar{\tilde{B}}_+\gamma_\mu\gamma_5 t^a\tilde{B}_+ + (c_2 - c_4)\bar{\tilde{B}}_-\gamma_\mu\gamma_5 t^a\tilde{B}_- \\ & + (1 - c_3)(\bar{\tilde{B}}_+\gamma_\mu t^a\tilde{B}_- + \text{h.c.}) + (\text{pion terms}) \end{aligned} \quad (18)$$

No predictions are obtained from chiral symmetry for these couplings, which can take any values.

The scenario of “effective chiral symmetry restoration” discussed by the authors of Ref. [6] requires that the coefficients of the chirally invariant operators m_1, c_{2-4} in this effective Lagrangian be suppressed. The alternative explanations for parity doubling described in the following subsections, would, in the language of $SU(2)_L \times SU(2)_R$ symmetry restoration, amount to a dynamical mechanism for the suppression of m_1 and c_{2-4} . For further discussion of parity doubling and $SU(2)_L \times SU(2)_R$ symmetry restoration, see Ref. [11]. Note that these results are specific to the assignment of $B_\pm$ to the chiral representations $(0, \frac{1}{2}), (\frac{1}{2}, 0)$, and a different chiral assignment will give different predictions for the pion couplings.

Finally, for completeness, we describe the alternative scenario suggested in 1969 by Dashen [32], and revisited recently by Kogan, Kovner, and Shifman[33], wherein the

vacuum is left invariant not only by the vector generators $\{T^a\}$, but also by a discrete symmetry $Z_{n_f}^A$ (the center symmetry of $SU(n_f)_A$). This scenario has no implications for parity doubling unless $n_f \geq 3$, so we restrict our consideration to the minimal case $n_f = 3$. Although at least three light flavors are required, this scenario would have nontrivial implications for the nonstrange baryons, which are our main focus here.




FIGURE 7. Typical baryon mass spectrum in the Dashen scenario for chiral symmetry breaking. Shown are two singlets of positive and negative parity, and a parity doublet. Pion transitions $h_1 \rightarrow h_2\pi$ between singlets and doublets are forbidden by the symmetry (dashed arrows), while transitions among singlets and among doublets are allowed (solid arrow).

The center of axial $SU(3)$ contains the operator $Z = \exp(\frac{2\pi i}{3}X_8)$ along with Z^2 and the identity. Since Z does not commute with parity, the complete discrete symmetry of the vacuum is the dihedral group $D_3 = (1, Z, Z^2, P, PZ, PZ^2)$. According to Coleman's theorem [18], the same symmetry should also be apparent in the hadron spectrum. Since D_3 has both one- and two-dimensional irreducible representations, the Dashen scenario of chiral symmetry realization could give rise to parity doubling in the hadron spectrum⁴. More precisely, there are one-dimensional representations of either parity, and two-dimensional representations, containing parity doublets.

Since mesons are singlets under the discrete D_3 group, this symmetry gives a selection rule for the pion strong decays between hadrons: the $h^* \rightarrow h\pi$ decay is allowed only between singlets, or only between doublets, but not between singlets and doublets as illustrated in Fig. 7. No clear evidence for this selection rule is observed among the baryons — in fact Dashen points out several problems with it that still pertain today — although symmetry breaking effects through mass terms could obscure it. We refer to the original work [32] for details.

In the Dashen scenario the $\langle \bar{q}q \rangle$ condensate vanishes in the massless quark limit, and $f_\pi \sim m_q$ as well. Phenomenological analysis of the pattern of chiral symmetry breaking [33, 34] does not rule out this behavior. However Kogan, Kovner, and Shifman have shown that it is in conflict with rigorous QCD inequalities [35]. So we do not consider the Dashen scenario further as a possible origin for parity doubling.

⁴ This does not occur in the two flavor case because $Z_2 \times \text{parity}$ is Abelian and has only one dimensional irreducible representations.

TABLE 1. Comparison of the physical manifestations of the scenario discussed here of “suppression of $U(1)_A$ symmetry breaking (SB)”, with those of “ $U(1)_A$ restoration”, which would be obtained by attempting to realize $U(1)_A$ in Wigner-Weyl mode.

Prediction	$U(1)_A$ restoration	Suppression of $U(1)_A$ SB
$Q_5 0\rangle = 0$	yes	no
parity doubling	yes	yes
mass states are Q_5 eigenstates	yes	no

Dynamical Suppression of $U(1)_A$ Symmetry Violation

In this section we adopt the standard picture of $U(1)_A$ symmetry violation in QCD as summarized, for example, by ’t Hooft[17]. The $U(1)_A$ symmetry is broken in the quantum Hamiltonian by quark masses and by the triangle anomaly. Since we are interested in baryons made of u , d , and s quarks, we consider QCD with three flavors. On account of the explicit symmetry breaking, the flavor singlet axial current, j_5^μ , has a divergence, and its charge, $Q_5(t) = \int d^3x j_5^0(t, \vec{x})$, does not commute with the Hamiltonian,

$$[H, Q_5] = i \frac{d}{dt} Q_5 = i \int d^3x \partial_\mu j_5^\mu, \quad \partial_\mu j_5^\mu = \sum_{q=u,d,s} 2m_q i \bar{q} \gamma_5 q - i \frac{3g^2}{16\pi^2} \text{Tr} \tilde{F}^{\mu\nu} F_{\mu\nu} \quad (19)$$

Because there are no massless particles coupled to j_5^μ , the singlet axial charge is a well defined operator. This contrasts with the isospin axial charges, Q_5^a , $a = 1, 2, 3$, which become ill-defined in the chiral limit.

Some of the consequences of $U(1)_A$ invariance, including parity doubling, can be obtained if the matrix elements of $[H, Q_5]$ are suppressed in a subspace of the space of states where the spectrum is sparse. The operator Q_5 can create η' mesons from the vacuum, so it clearly contains frequency components of order 1 GeV. We believe this obstructs restoration of $U(1)_A$ symmetry in a *strong*, operator, sense. However a *weak* form, applied only to matrix elements within the subspace, seems possible and could lead to parity doubling.

Suppose first, for simplicity, that there were a subspace, $\mathfrak{h}$, of the Hilbert space of QCD, in which the matrix elements of $\partial_\mu j_5^\mu$ were exactly zero. One possibility is that Q_5 also has no non-zero matrix elements in this subspace. In that case the symmetry has no implications for the spectrum in $\mathfrak{h}$. Alternatively, suppose Q_5 has a non-zero matrix element between two mass eigenstates $|B_+\rangle$ and $|B_-\rangle$ in $\mathfrak{h}$. B_+ and B_- must have the same flavor and angular momentum quantum numbers but opposite parity, denoted by the $\pm$ sign. According to eq. (19) they must also have the same mass if $\partial_\mu j_5^\mu$ can be neglected as we have assumed,

$$\langle B_+ | [Q_5, H] | B_- \rangle = (m(B_-) - m(B_+)) \langle B_+ | Q_5 | B_- \rangle = 0 \quad (20)$$

If $\langle B_+ | Q_5 | B_- \rangle \neq 0$, as we have assumed, then $m(B_+) = m(B_-)$. So the spectrum of states with a given J , I , and S , consists either of isolated states which have no non-zero matrix

elements of Q_5 within $\mathfrak{h}$, or of degenerate families of both positive and negative parity connected by Q_5 . Excluding accidental degeneracies, the result is parity doubling.

We do not believe that stronger conditions can be imposed:

- a. $[H, Q_5] = 0$ and $Q_5|0\rangle = 0$

This condition is not consistent with the QCD lagrangian. Phenomenologically, it leads to parity doubling throughout the baryon spectrum. Since parity doubling is at best an approximate symmetry in a subspace, and clearly does not hold for the lightest states, this condition is not tenable.

- b. $[H, Q_5]|B\rangle = 0$ if $B \in \mathfrak{h}$

It is easy to show that this condition leads to parity doubling of the state $|B\rangle$, however we believe it is too strong. In particular, it implies that the state $Q_5|B\rangle$ must be an eigenstate of H with the same mass as $|B\rangle$. We expect that the operator Q_5 generically creates η' mesons, so the amplitude $\langle B'\eta'|Q_5|B\rangle$, where $|B'\rangle$ might be the original state $|B\rangle$ or some other baryon, should not be small. The weak condition we have adopted does not preclude the possibility that the axial charge acting on one member of a parity doublet has a significant amplitude to produce an η' .

If the matrix elements of $\partial_\mu j_5^\mu$ are small, but not zero, in $\mathfrak{h}$, then the degeneracies are no longer exact. It is interesting to consider this case from the point of view of the $U(1)_A$ Goldberger-Treiman relation and to contrast it with the $SU(2)_L \times SU(2)_R$ case.⁵ Again taking $B_\pm$ to be spin-1/2 baryons, the matrix element of the flavor singlet axial current and its divergence can be parameterized in terms of invariant form factors,

$$\begin{aligned}\langle B_+|j_{5\mu}(0)|B_- \rangle &= \bar{u}(p_+) (\gamma_\mu g_A(q^2) + q_\mu g_P(q^2) + i\sigma_{\mu\nu} q^\nu g_M(q^2)) u(p_-) \\ \langle B_+|\partial^\mu j_{5\mu}|B_- \rangle &= \bar{u}(p_+) d_A(q^2) u(p_-)\end{aligned}\tag{21}$$

Taking the states to be at rest, $q^2 = (q^0)^2 = \Delta m^2$. Then contracting q^μ with the first of eqs. (21) we obtain

$$\Delta m (g_A(\Delta m^2) + \Delta m g_P(\Delta m^2)) = d_A(\Delta m^2)\tag{22}$$

Note that the quantity in parentheses is the matrix element of the axial charge: $\langle B_+|Q_5|B_- \rangle \propto g_A(\Delta m^2) + \Delta m g_P(\Delta m^2)$. The flavor singlet case differs from the flavor non-singlet case in that there is no zero mass pole in the induced pseudoscalar form factor, g_P . Indeed, the nearest significant singularity in the pseudoscalar channel is the η' at nearly 1 GeV. Therefore, if the matrix elements of $\partial_\mu j_5^\mu$ are small, the induced pseudoscalar contribution is negligible (second order in Δm^2), leaving,

$$\Delta m g_A(0) \approx d_A(0)\tag{23}$$

Eq. (23) allows us to formulate the problem of “suppression of $U(1)_A$ symmetry violation”: Are there states for which $d_A(0)$ is small and $g_A(0)$ is not? While we cannot answer the question decisively, it is well posed. At least in principle, this question can be

⁵ The $U(1)_A$ Goldberger-Treiman relation was considered for nucleons in [36].

answered using lattice QCD computations of the form factor $d_A(q^2)$. In the chiral limit, this coincides with the form factor of the gluonic term in the anomaly.

We emphasize that the scenario of “suppression of $U(1)_A$ symmetry violation” is very different from that of $U(1)_A$ restoration, or $U(1)_A$ realized in Wigner-Weyl mode. For convenience we summarize in Table 1 the similarities and differences between these two scenarios. In particular, we do not require the invariance of the vacuum under $U(1)_A$ transformations.

Quark models suggest that the matrix elements of the singlet axial current between states with similar quark configurations but opposite parity are of order unity. For example the singlet axial current in the bag model changes a quark with angular momentum j and positive parity into a quark with angular momentum j and negative parity with amplitude of order unity. For such states $\Delta m \sim d_A(0)$, and suppression of the matrix elements of the divergence of the singlet axial current would result in approximate parity doubling. Other states, for which $g_A(0)$ is smaller, need not be nearly degenerate, even though $d_A(0)$ may be small.

The divergence of the flavor singlet axial current gets contributions from the anomaly and from quark masses. Is it possible that the matrix elements of $\tilde{F}F$ and $m_s \bar{s} \gamma_5 s$ are unusually small at moderate excitations in the baryon spectrum? Perhaps instanton fluctuations are suppressed at high excitation in the baryon spectrum as they are at short distances. We are not aware of any work along this direction.⁶ We do not know how to estimate the matrix elements of $F\tilde{F}$ between excited hadrons, and leave it for further study.

The generalization of suppression of $U(1)_A$ symmetry violation to strange hadrons is not straight forward. In the limit $m_s = 0$ $SU(3)_V$ becomes exact and therefore suppression of $\langle F\tilde{F} \rangle$ would lead to parity doubling among strange as well as non-strange baryons in octets and decuplets. When the strange quark mass is increased, the flavor singlet current has an increasing divergence proportional to $m_s \bar{s} \gamma_5 s$. However there is a linear combination of flavor singlet and non-singlet currents that includes only u and d quarks, $J_5^\mu \equiv \bar{u} \gamma^\mu \gamma_5 u + \bar{d} \gamma^\mu \gamma_5 d$, whose divergence comes only from the axial anomaly,

$$\partial_\mu J_5^\mu = -i \frac{2g^2}{16\pi^2} \text{Tr} \tilde{F}^{\mu\nu} F_{\mu\nu}$$

The discussion of parity doubling among non-strange hadrons can be repeated without modification using J_5^μ instead of the flavor singlet current, j_5^μ . However $\langle N^\pm, \Delta^\pm | F\tilde{F} | N^\mp, \Delta^\mp \rangle \approx 0$ does not imply that $F\tilde{F} \approx 0$ for their strange $SU(3)$ partners. For example, one might model the effects of the anomaly by the 't Hooft determinant, $\det \bar{q}_R q_L + \det \bar{q}_L q_R$, which is a six-quark operator of the form $\bar{u} \bar{d} \bar{s} u d s$ [41]. This operator has tree level matrix elements in Λ 's and Σ 's, but only contracted loop contributions in non-strange baryons. Without a better model of hadron matrix elements of $F\tilde{F}$, it is not

⁶ In Ref. [37] Shifman's analysis [8] is extended to flavor singlet correlators, where instanton contributions determine the convergence of sum rules analogous to Weinberg's [40]. Our analysis suggests that the proper quantity to analyze would be matrix elements of $\tilde{F}F$ between baryon states of identical J , I , and S , but opposite parity, possibly using instanton methods along the lines of ref. [38].

possible to conclude that parity doubling among non-strange baryons implies doubling among the analogous strange states.

Finally it is worth emphasizing that this sort of symmetry restoration occurs frequently in quantum systems. Since the phenomenon involves *explicit* rather than *spontaneous* symmetry violation, we can find examples in ordinary quantum mechanics. Consider, for example, an atom in an external electric quadrupole field, $\delta H_{SB} \propto e(r_i r_j - \frac{1}{3}\delta_{ij})Q^{ij}$. Although rotational invariance is broken in general, it is unbroken among $j = 1/2$ states where $\langle \psi_{\frac{1}{2},m} | [\delta H_{SB}, \vec{J}] | \psi'_{\frac{1}{2},m'} \rangle = 0$ even though $[\delta H_{SB}, \vec{J}] | \psi_{\frac{1}{2},m} \rangle \neq 0$ in general.

Parity Doubling and Chiral Symmetry Restoration at Short Distances

The idea that $SU(2)_L \times SU(2)_R$ chiral symmetry is restored at asymptotically large Euclidean momenta (short distances) in current correlators in QCD dates back to Weinberg's 1967 paper on spectral function sum rules. These sum rules relate moments of the spectral functions of vector and axial currents. However, turning them into relations among individual meson states is a different problem, and requires additional theoretical input.

Recently, such relations have been discussed by several authors [8, 47], who argued that parity doubling might be a consequence of the chiral invariance of current correlators at short distances in QCD and other dynamical assumptions. Usually these arguments are framed in terms of “chiral symmetry restoration”, so it is interesting to ask, in the spirit of Ref. [11], what additional dynamics are responsible for suppressing the chirally invariant operators necessary to obtain a Wigner-Weyl realization of chiral symmetry in the spectrum.

Refs. [8, 47] discuss mesons at high excitation and large N_c . The arguments do not go over to the baryon sector unchanged [48] since baryons have different large N_c systematics than mesons, and baryon correlators have different short distance behavior than meson current correlators. Furthermore our phenomenological analysis pertains to hadrons in the 1.0 — 2.5 GeV region, and includes many resonances that are the lightest states in a particular isospin and angular momentum channel, where such an analysis certainly would not apply. Nevertheless it remains interesting to examine the assumptions required to obtain parity doubling in this simpler limiting case. Ref. [47] discusses relations among vector and axial mesons, while Ref. [8] presents similar relations for pseudoscalar and scalar mesons. For simplicity, we phrase our discussion in terms of the latter case.

Shifman compares the correlators of two scalar currents, $\Pi(Q^2)$, with the correlator of two pseudoscalar currents, $\tilde{\Pi}(Q^2)$, in the Euclidean domain, $Q^2 = -q^2 > 0$. Shifman assumes that the difference, $\Delta\Pi(Q^2) \equiv \Pi(Q^2) - \tilde{\Pi}(Q^2)$ obeys an unsubtracted dispersion relation,

$$\Delta\Pi(Q^2) \equiv \Pi(Q^2) - \tilde{\Pi}(Q^2) = - \int d\sigma^2 \frac{\Delta\rho(\sigma^2)}{Q^2 + \sigma^2} \quad (24)$$

where the spectral functions $\rho(\sigma^2)$ and $\tilde{\rho}(\sigma^2)$ (with $\Delta\rho \equiv \rho - \tilde{\rho}$) receive contributions from physical intermediate states of mass σ created by the currents from the vacuum. Well established arguments based on the operator product expansion and QCD sum

rules[49] require that $\Delta\Pi(Q^2)$ vanishes like $1/Q^4$ as $Q^2 \rightarrow \infty$,

$$\Delta\Pi(Q^2) \underset{Q^2 \rightarrow \infty}{\sim} \frac{\langle \bar{\Psi}\Psi \rangle^2}{Q^4} \quad (25)$$

where $\langle \bar{\Psi}\Psi \rangle$ is the chiral symmetry violating condensate. Each of the correlators, Π and $\tilde{\Pi}$, individually go like $Q^2 \log Q^2$ as $Q^2 \rightarrow \infty$, so the fact that their difference vanishes as $1/Q^4$ is evidence of the rapid restoration of chiral symmetry *at short distances* in QCD. Combined with the unsubtracted dispersion relation, the condition that there is no term $\sim 1/Q^2$ in $\Delta\Pi(Q^2)$ requires a superconvergence relation,

$$\int d\sigma^2 \Delta\rho(\sigma^2) = 0 \quad (26)$$

Eq. (26) is the *only* input from chiral symmetry in the argument of Ref. [8]. Convergence of the σ integration requires that the spectral functions, each of which grows like σ^2 must quickly become equal at large σ . Naively one would think that this constraint applies at asymptotically large masses where broad, overlapping resonances build up continua in ρ and $\tilde{\rho}$. Strong additional assumptions are needed to make inferences about parity doubling. The principal assumption is that QCD may be described by a limit in which the number of colors, N_c , is taken to infinity *before* considering high radial excitations (labeled by a radial quantum number n).

To explore the implications of this unusual order of limits, we analyze the problem at large but fixed N_c and extract the leading order predictions at the end. First, let us recap the argument of Ref. [8], taking $N_c \rightarrow \infty$ first. Then, the spectral function ρ ($\tilde{\rho}$) receives contributions only from *zero width* scalar (pseudoscalar) meson resonances, with masses M_n ($\tilde{M}_n$) and coupling to the current f_n ($\tilde{f}_n$). Continuum contributions are suppressed by powers of N_c . For reference, the squared masses, squared widths, and couplings of the mesons are assumed to have the following dependence on n and N_c

$$\begin{aligned} M_n^2, \tilde{M}_n^2 &\sim \Lambda^2 n \\ \Gamma_n^2, \tilde{\Gamma}_n^2 &\sim \Lambda^2 n / N_c^2 \\ f_n, \tilde{f}_n &\sim N_c \Lambda^4 n \end{aligned} \quad (27)$$

where Λ is the dynamical mass scale of QCD. The n dependence is motivated by string analogies, the spectra of linear potentials and the 't Hooft model[8, 51]. The factors of N_c are in accord with standard large- N_c counting arguments.

Ignoring the widths or equivalently, exchanging the limit $N_c \rightarrow \infty$ with the sum over n , the spectral functions are sums of poles on the real axis, so

$$\Delta\rho(\sigma^2) = \sum_n (f_n \delta(\sigma^2 - M_n^2) - \tilde{f}_n \delta(\sigma^2 - \tilde{M}_n^2)) \quad (28)$$

and the superconvergence relation reads,

$$N_c \Lambda^2 \sum_n (\alpha_n M_n^2 - \tilde{\alpha}_n \tilde{M}_n^2) = 0 \quad (29)$$

where we have defined $f_n = N_c \Lambda^2 \alpha_n$, etc. This sum must converge at large n . Taking $\alpha_n = \tilde{\alpha}_n = \alpha$, independent of n (an assumption later relaxed in Ref. [8]) the weakest behavior large n behavior consistent with the convergence of the sum in eq. (29) is

$$M_n^2 - \tilde{M}_n^2 \underset{n \rightarrow \infty}{\sim} \text{alt} \frac{1}{n}$$

where “alt” denotes an alternating sign. Since $M_n \sim \tilde{M}_n \sim \sqrt{n}$ at large n , Shifman finds $\Delta M_n \equiv M_n - \tilde{M}_n \sim \text{alt} n^{-3/2}$, indicating that the spectrum achieves parity doubling rather rapidly at large n ⁷.

For any fixed N_c the widths of resonances grow at the same rate as their masses (see eq. (27)), so the spectral functions approximate a continuum as $n \rightarrow \infty$ at fixed N_c . Let us keep track of these $1/N_c$ corrections. The effect of a finite width for the n^{th} meson on the spectral functions was analyzed in Ref. [52]. When the widths are finite the meson poles in the correlation functions move into the lower half complex q^2 -plane, and the correlation functions develop cuts with branch points at the thresholds for the decay of the n^{th} meson. We refer to Ref. [52] for details. It is straightforward to find the modification of eq. (29)

$$N_c \Lambda^2 \sum_n \left(\alpha_n (M_n^2 - \frac{1}{\pi} \Gamma_n M_n) - \tilde{\alpha}_n (\tilde{M}_n^2 - \frac{1}{\pi} \tilde{\Gamma}_n \tilde{M}_n) \right) = 0 \quad (30)$$

Repeating the same arguments that followed eq. (29), we find

$$\Delta M_n - \frac{1}{\pi} \Delta \Gamma_n \underset{n \rightarrow \infty}{\sim} \text{alt} \frac{1}{n^{3/2}}$$

where $\Delta \Gamma_n \equiv \Gamma_n - \tilde{\Gamma}_n$. Referring back to eqs. (27) we see that $\Gamma, \tilde{\Gamma} \sim \sqrt{n}/N_c$ to leading order in n and N_c . However we have no independent argument to fix the large n behavior of the difference $\Delta \Gamma_n$. It could, in principle go like a constant at large n , or like some power of n larger than $n^{-3/2}$. If we parameterize our ignorance by an exponent β , presumably smaller than $1/2$, we can summarize the $1/N_c$ corrections to the analysis of Ref. [8],

$$\Delta M_n \underset{n \rightarrow \infty}{\sim} \text{alt} \frac{1}{n^{3/2}} + C \frac{n^\beta}{N_c}$$

where both β and the constant C are unknown. Thus the effect of $1/N_c$ (remember $N_c = 3$ in our world) corrections could completely destroy parity doubling. While further

⁷ This derivation has been criticized in Ref. [50], where it is pointed out that the $N_c \rightarrow \infty$ correlator, $\Delta \Pi(Q^2)$, does not satisfy an unsubtracted dispersion relation, or equivalently that certain interchanges of limits and sums/integrals, required by the derivation, are not allowed by the asymptotic behavior of $\Delta \rho(\sigma^2)$ when N_c is infinite. By keeping finite N_c corrections (see the following paragraph), which replace the δ -functions in $\Delta \rho$ with finite width resonances (in the manner proposed in Ref. [52]), we maintain the unsubtracted superconvergence relation and avoid any problems with interchanges of limits. By this approach we include important physics, namely the fact that the resonant contributions to the spectral functions merge into a continuum at large n for any finite N_c , and, happily, agree with the conclusions of Ref. [50] that asymptotic chiral symmetry and conventional large N_c arguments alone do not require parity doubling.

analysis or more assumptions might indicate that these $1/N_c$ effects can be neglected, this example serves well to illustrate the fact that parity doubling does not follow from the chiral invariance of QCD alone, nor from suppression of chiral symmetry *violating* effects. In the example of Ref. [8] the assumptions include those necessary to give the asymptotic behavior summarized in eqs. (27) and to suppress the effects of hadron widths at large but finite N_c .

Parity Doubling and Intrinsic Deformation

Parity doubling in the baryon spectrum might be a consequence of a phenomenon well known in nuclear and molecular physics. If a system can be described by the collective quantization of an intrinsic state and a) the intrinsic state (spontaneously) violates parity, and b) the deformation is relatively rigid, then the low lying excitations of the system will display parity doubling. The phenomenon is well known, for example, in the spectrum of ammonia. Clear examples occur in nuclear physics as well. The existence of parity doubled states in nuclei like ^{225}Ra , with intrinsic octupole (“pear shaped”) deformation, plays an important role in proposed sensitive searches for T -violating electric dipole moments[42, 43, 44]. Linear superpositions of the intrinsic state, $|\psi\rangle$ and its parity image, $\hat{P}|\psi\rangle$ form parity eigenstates, $|\psi_{\pm}\rangle = \frac{1}{\sqrt{2}}(|\psi\rangle \pm \hat{P}|\psi\rangle)$. If the deformation is rigid then the Hamiltonian matrix element between $|\psi\rangle$ and $\hat{P}|\psi\rangle$ is small and the states of opposite parity, but otherwise identical structure, are nearly degenerate. This degeneracy survives quantization of the collective coordinates that restores rotational symmetry resulting in a tower of parity doubled excitations.

One can imagine dynamics that might lead to a similar phenomenon in hadron spectroscopy [2], though a convincing discussion is beyond our present understanding of hadrons in QCD. If baryons on the leading Regge trajectory were well described as a dumbbell-like structure with a quark on one end and a diquark on the other[45], then the intrinsic state would violate reflection symmetry. If tunneling of the odd quark from one end to the other were suppressed enough, then parity doubling would result. If correct, this argument would predict parity doubling for states of high angular momentum. There is some evidence for this among nucleon ($I = 1/2$) and Δ ($I = 3/2$) states, although the Δ s are not very well known (see Table 2). Hyperons (Λ s and Σ s) are too poorly known to provide much information. If this mechanism is responsible for parity doubling, one expects the evidence for parity doubling should improve with increasing angular momentum, and should be found in the spectrum of Λ s and Σ s as well. These predictions could be tested in an experimental re-examination of baryon spectroscopy. As a first attempt in this direction, we have repeated the analysis of Section 2 on the non-strange baryons with the states in Table 2 *removed*. The significance of the signal for parity doubling remains virtually unchanged when the high angular momentum states are removed. Thus the phenomenon cannot be entirely explained by collective coordinate quantization of a parity-violating, elongated intrinsic state.

TABLE 2. Parity doubled $I = 1/2$ and $I = 3/2$ baryons with angular momentum $5/2$ or greater. Data from the PDG[29].

$B(J^P)$	PDG reliability	Mass (MeV)
$N(5/2^-)$	****	1675
$N(5/2^+)$	****	1680
$N(7/2^+)$	**	1990
$N(7/2^-)$	****	2190
$N(9/2^+)$	****	2220
$N(9/2^-)$	****	2250
$\Delta(5/2^+)$	****	1905
$\Delta(5/2^-)$	***	1930
$\Delta(7/2^+)$	****	1950
$\Delta(7/2^-)$	*	2200
$\Delta(9/2^+)$	**	2300
$\Delta(9/2^-)$	**	2400

IMPLICATIONS FROM MENDED CHIRAL SYMMETRY

We explore in this Section some implications of chiral symmetry using a comparatively less well-known approach. While not directly relevant to the topic of parity doubling, this approach has potentially interesting implications for the properties of baryon resonances of opposite parity. As discussed in Section 2, chiral symmetry realized in Nambu-Goldstone mode does not impose any constraints on the masses and couplings of hadrons, beyond those required by isospin. However, in a series of interesting papers, Weinberg showed that the full chiral symmetry group may remain manifest in a very special sense [23, 24]. The symmetry is realized on the matrix of the couplings of the Goldstone bosons, rather than on the mass eigenstates, as typical for a symmetry realized in Wigner-Weyl mode. Following Ref. [24], we will refer to chiral symmetry realized in this fashion as “mended chiral symmetry”. When combined with the large energy asymptotics of Goldstone boson scattering amplitudes following from Regge theory, the approach of Refs. [23, 24] gives also nontrivial implications for the mass spectrum.

In this section we give a brief review of the approach of Refs. [23, 24], and use it to construct a chirally invariant model for a system of two nucleons of opposite parity. In this model the two nucleons of opposite parity are not necessarily degenerate, but their pion couplings are exactly the same as predicted from a linear realization of chiral symmetry. This approach provides such a natural realization of such predictions of chiral symmetry “restoration” without actual parity doubling.

The mended chiral symmetry approach is equally applicable to mesons and baryons. In the original work Ref. [23] it was used to describe the quartet of $J^P = 0^\pm, 1^\pm$ mesons (σ, π, ρ, a_1), and the (N, Δ, N_1) system of the nucleon, delta and Roper baryons, respectively. However, in the sector of mesons with zero helicity, this approach is considerably more predictive than in the baryon sector with nonzero helicity. This is due to the existence of an endomorphism following from the parity transformation of the

pion couplings. We will restrict our discussion in this section to the predictions which are common to both baryons and mesons.

We start by introducing some notation. For simplicity, we consider $n_f = 2$ with isospin generators T^a . Consider the amplitude $\mathcal{M}_{\alpha\beta}^a$ for pion coupling between two states $h_\alpha(\lambda) \rightarrow h_\beta(\lambda')\pi^a$ with h_α, h_β baryon states with helicities λ, λ' , respectively.

This amplitude is related to the matrix element of the axial vector current between the two baryon states by PCAC as

$$A(h_\alpha(p, \lambda) \rightarrow h_\beta(p', \lambda')\pi^a) \equiv \mathcal{M}_{\alpha\beta}^a = \frac{1}{f_\pi}(p - p')_\mu \langle h_\beta(p', \lambda') | A^{a\mu} | h_\alpha(p, \lambda) \rangle \quad (31)$$

The form of the pion coupling simplifies in the collinear limit, where the baryon momenta are parallel to a common direction $\vec{p} \parallel \vec{p}' \parallel \hat{e}_3$, which can be chosen for convenience to define the $+z$ direction. In such a frame, the pion coupling given by the expression Eq. (31) is related to the matrix element of a light-cone component of the axial current, and is helicity conserving

$$\mathcal{M}_{\alpha\beta}^a = \frac{1}{f_\pi} |\vec{q}| \langle h_\beta(p', \lambda') | A^{a0} - A^{a3} | h_\alpha(p, \lambda) \rangle \equiv [X^a(\lambda)]_{\beta\alpha} \delta_{\lambda\lambda'} \quad (32)$$

This defines the transition operator $X^a(\lambda)$ for pion couplings to hadrons. For simplicity we will omit the λ dependence of the operators $X^a(\lambda)$, unless explicitly required.

The pion couplings X^a and the isospin operators T^a satisfy the commutation relations

$$[T^a, T^b] = i\epsilon^{abc} T^c, \quad [T^a, X^b(\lambda)] = i\epsilon^{abc} X^c(\lambda). \quad (33)$$

In addition to this, another commutation relation can be obtained by considering the set of all Adler-Weisberger sum rules [53] on baryon targets. They can be derived from an unsubtracted dispersion relation for the $h_\alpha\pi^a \rightarrow h_\beta\pi^b$ scattering amplitude. Saturating the dispersion relation with one-body intermediate states, the sum rule can be expressed in operator form as

$$[X^a(\lambda), X^b(\lambda)] = i\epsilon^{abc} T^c \quad (34)$$

The commutation relations Eqs. (33), (34) form the Lie algebra of the $SU(2)_L \times SU(2)_R$ symmetry, which is furthermore explicitly realized on the hadronic states. This realization may be different for each distinct helicity λ . The resonance saturation approximation of the Adler-Weisberger sum rule, required to derive the commutation relation Eq. (34), can be justified, for example, by working at leading order in the $1/N_c$ expansion.

The absence of the $\Delta I = 2$ Regge trajectories with intersect $\alpha(0) > 0$ for the pion-nucleon scattering amplitude $h_\alpha\pi \rightarrow h_\beta\pi$ gives another commutation relation. This connects the pion coupling operator X^a with the mass spectrum

$$[X^a(\lambda), [X^b(\lambda), m^2]] \propto \delta_{ab} \quad (35)$$

This commutation relation implies that the baryon mass operator has very simple transformation properties under $SU(2)_L \times SU(2)_R$. It consists of two terms

$$\hat{m}^2(\lambda) = \hat{m}_0^2(\lambda) + \hat{m}_4^2(\lambda) \quad (36)$$

with $\hat{m}_0^2$ transforming as a singlet $(0_L, 0_R)$ under the chiral group $SU(2)_L \times SU(2)_R$, and $\hat{m}_4^2$ transforming as the $I = 0$ component of a chiral four-vector $(\frac{1}{2}L, \frac{1}{2}R)$. In the absence of the $\hat{m}_4^2$ term, the chiral symmetry would have been realized explicitly in the mass spectrum, and baryons would be grouped into irreducible representations of $SU(2)_L \times SU(2)_R$. The $\hat{m}_4^2$ term prevents such a simple picture, and implies that in general, mass eigenstates are mixtures of different irreducible representations of $SU(2)_L \times SU(2)_R$ and do not exhibit the pattern of degeneracies associated with a Wigner-Weyl representation of $SU(2)_L \times SU(2)_R$. Note that $\hat{m}_4^2$ is *not* connected with explicit chiral symmetry breaking, which has been set to zero from the outset. Thus Weinberg's mended symmetry scenario provides a good example of the point emphasized in Ref. [11], that "restoration" of $SU(2)_L \times SU(2)_R$ in the spectrum requires suppression of chirally invariant operators.

To summarize, Weinberg's dynamical arguments, based on assumed Regge asymptotics, suffice to prove that hadrons with given helicity λ fall into *reducible* representations of the chiral group. These representations in general may be different for different helicities. The reducible nature of the representations and the absence of degeneracies distinguish Weinberg's approach from the "chiral symmetry restoration" picture discussed in Sec. III. The crucial feature is the presence of the chiral nonsinglet $\hat{m}_4^2$ piece in the Hamiltonian, which connects different irreps, according to selection rules imposed by its quantum numbers. The representation content of a given hadron state is not fixed by these general arguments.

The mended symmetry approach, in effect, replaces the arbitrariness of the coupling constants and masses in the effective Lagrangian with the equally unknown representation content of a given state. This framework is predictive only when supplemented with a prescription for the chiral irreps content. Once the representation content is specified, many physical parameters (masses and couplings) are predicted in terms of just a few input parameters. This lends such an approach considerable predictive power. For applications of this formalism to heavy hadron physics and exotics see Ref. [56, 57, 58, 59].

We comment here on the relation of this approach to the large N_c picture of the baryons following from the work of Dashen, Jenkins and Manohar [25, 26]. In the large N_c limit, a new symmetry emerges in the baryon sector of QCD. This is the contracted spin-flavor symmetry $SU(2n_f)_c$, with n_f the number of light quark flavors. The generators of the symmetry are the spin J^i , isospin I^a , and the operator G_0^{ia} defined in terms of p-wave pion couplings to the baryons in the large N_c limit ($G^{ia} \rightarrow G_0^{ia}$)

$$\mathcal{M}_{\alpha\beta}^a = \frac{1}{f_\pi} q^i N_c [G^{ia}]_{\beta\alpha} \quad (37)$$

The connection to the operators introduced in Eq. (32) is $X^a(\lambda) \sim [G^{3a}]_{\lambda\lambda}$.

The operators J^i, T^a, G_0^{ia} satisfy the commutation relations defining the algebra of the contracted $SU(2n_f)$ symmetry

$$[J^i, J^j] = i \varepsilon_{ijk} J^k, \quad [T^a, T^b] = i \varepsilon_{abc} T^c \quad (38)$$

$$[G_0^{ia}, J^j] = i \varepsilon_{ijk} G_0^{ka}, \quad [G_0^{ia}, T^b] = i \varepsilon_{abc} G_0^{ic}, \quad [G_0^{ia}, G_0^{jb}] = 0. \quad (39)$$

Furthermore, the mass operator in the large N_c limit m_0^2 commutes with all the generators of the contracted symmetry.

The contracted $SU(2n_f)$ symmetry relates states with different spins and isospins, analogous to the Wigner symmetry in nuclear physics. For example, in the symmetry limit the nucleon N and the Δ belong to the same irreducible representation of the $SU(4)_c$, along with infinitely many other states with spin and isospin $J = I = 1/2, 3/2, 5/2, \dots$. The pion couplings among these states are fixed by the contracted symmetry, such that they are precisely the same as the predictions of the constituent quark model [25, 26].

Both the mended chiral symmetry and the large N_c contracted symmetry approach impose constraints on the form of the pion couplings to the baryons. In general these constraints are different, although they can agree in special situations. The reason for the difference is the commutation relation Eq. (35). In the large N_c limit such a commutator would vanish, since the mass operator commutes with the pion couplings $X^a(\lambda)$. This leads to a nontrivial chiral structure of the mass operator in the mended symmetry approach, which does not follow from the large N_c contracted symmetry.

In certain particular cases this difference becomes immaterial, and the predictions of the two approaches are identical. This happens when the hadrons in a sector of given helicity λ are assigned to one single irrep (I_L, I_R) of $SU(2)_L \times SU(2)_R$, which renders the double commutator Eq. (35) zero, in agreement with the large N_c prediction. Such a situation is illustrated in Scenario I below.

Another important difference between the two approaches lies in their range of applicability: while the mended chiral symmetry method is restricted to the couplings of the Goldstone bosons to baryons, the contracted large N_c symmetry is equally applicable to the couplings of any light mesons to baryons.

We proceed to an explicit discussion of the constraints on the mass spectrum and pion couplings from the mended chiral symmetry approach. The hadronic states with given helicity λ can be taken to transform as linear combinations of irreducible representations (j_L, j_R) of $SU(2)_L \times SU(2)_R$

$$|h(\lambda)\rangle \sim \sum_{j_L, j_R} c_{j_L j_R}(\lambda) |j_L, j_R\rangle \quad (40)$$

where we have suppressed all labels on the states except for their $SU(2)_R \times SU(2)_L$ representation content. In general, the mass operator $\hat{m}_4^2$ will mix different representations (barring accidental degeneracies), such that the states $|h(\lambda)\rangle$ will be nondegenerate, unless the sum over representations contains only one term.

Parity invariance relates the representation contents of hadronic states with opposite helicity.⁸ Acting with the parity operator $\hat{P}$ on a hadron state $h_\alpha(\vec{p}, \lambda)$ gives a state of opposite helicity and momentum $\hat{P}|h_\alpha(\vec{p}, \lambda)\rangle = \pi_h |h_\alpha(-\vec{p}, -\lambda)\rangle$, with π_h the intrinsic parity of the state h . It is more convenient to consider the *normality* operator $\Pi = \hat{P}\hat{R}_\pi^\gamma$, the product of parity and a 180° rotation, which restores the momenta to their original direction. The action of this operator on the hadron states is $\Pi|h_\alpha(\vec{p}, \lambda)\rangle = \pi_h(-)^{j_h} |h_\alpha(\vec{p}, -\lambda)\rangle$.

⁸ Of course Lorentz invariance constrains the representation content so that states of the same hadron with different $|\lambda|$ are degenerate.




FIGURE 8. Weight diagrams for the chiral representations $(1, \frac{1}{2})$ and $(\frac{1}{2}, 1)$ for states with helicity $\lambda = +\frac{1}{2}$ (left) and $\lambda = -\frac{1}{2}$ (right), respectively. The states on the same diagonal have the same third component of the isospin $I^3 = \frac{1}{2}(I_L^3 + I_R^3)$, and correspond to the nucleons and Δ states shown.

The generators of the symmetry transform under Π as

$$\Pi X^a(\lambda)\Pi = -X^a(-\lambda), \quad \Pi T^a\Pi = T^a \quad (41)$$

The mass operator commutes with the normality operator. Using this relation, it is easy to see that the states $|h(-\lambda)\rangle$ furnish also a reducible representation of the chiral algebra, containing the representations (j_R, j_L) obtained from those in Eq. (40) by exchanging $j_L \leftrightarrow j_R$

$$|h(-\lambda)\rangle \sim \sum_{j_L, j_R} c_{j_R j_L}(\lambda) |j_L, j_R\rangle \quad (42)$$

To illustrate the application of this approach, we consider two particular cases. The first scenario describes a N, Δ pair with the same mass, which reproduces the $NN\pi, N\Delta\pi, \Delta\Delta\pi$ couplings of the quark model. The second scenario describes a pair of spin 1/2 nucleons of opposite parity.

Scenario I. A model for a nonstrange nucleon and delta baryons can be constructed by placing the hadrons into minimal chiral multiplets. This model was first proposed in Ref. [23] and discarded because it has the unphysical feature that the nucleon and delta are degenerate. It was proposed again in Ref. [55] as a possible justification for the quark model with $SU(4)$ spin-flavor symmetry. This scenario comes closest to the large N_c picture of the ground state baryons, where the nucleon- Δ mass splitting is of order $1/N_c$.

We place each helicity sector into irreducible representations of $SU(2)_L \times SU(2)_R$

$$\lambda = +\frac{1}{2} : (1_L, \frac{1}{2}_R) \supset N, \Delta \quad (43)$$

$$\lambda = +\frac{3}{2} : (\frac{3}{2}_L, 0_R) \supset \Delta \quad (44)$$

The transformation properties of the negative helicity states are obtained from these via the parity relation Eq. (42) and are

$$\lambda = -\frac{1}{2} : \left(\frac{1}{2}_L, 1_R\right) \supset N, \Delta \quad (45)$$

$$\lambda = -\frac{3}{2} : \left(0_L, \frac{3}{2}_R\right) \supset \Delta \quad (46)$$

We show in Figure 8 the weight diagrams of the chiral representations corresponding to $\lambda = \pm\frac{1}{2}$, and the associated hadron states.

The mass matrix (see eq. (36)) for the helicity $\lambda = +1/2$ is proportional to the unit matrix in the basis $(N_{+1/2}, \Delta_{+1/2})$, since the $\hat{m}_4^2$ term does not have diagonal matrix elements on an irreducible representation (I_L, I_R) . Thus the nucleon and delta are degenerate in this model.

Collinear pion emission connects only states with the same helicity, and the corresponding axial matrix elements can be obtained by expressing the axial current in terms of the generators of the chiral group $X^a = \frac{1}{2}(I_R^a - I_L^a)$. Defining the reduced matrix elements for p-wave pion emission as

$$\langle J'I'; \lambda I'_3 | X^a | JI; \lambda I_3 \rangle = \frac{1}{2J'+1} X(J', J) \langle J'\lambda | J1; \lambda 0 \rangle \langle I'I'_3 | I1; I_3 a \rangle \quad (47)$$

one finds $X(1/2, 1/2) = -5/\sqrt{2}$, $X(1/2, 3/2) = -4$, $X(3/2, 3/2) = -5\sqrt{2}$, in agreement with the predictions of the quark model with SU(4) spin-flavor Wigner symmetry. In particular, the result for $X(1/2, 1/2)$ is equivalent to $g_A/g_V = 5/3$. The agreement with the experimental result $g_A/g_V = 1.2695 \pm 0.0029$ can be improved by refining the model and adding a second chiral representation $(\frac{1}{2}_L, 0_R)$ in the $\lambda = +\frac{1}{2}$ sector. The mixing angle of the two representations is chosen appropriately such as to reproduce the data on g_A/g_V [23]. We do not consider this further, and refer the reader to Weinberg's paper for details [23].

Scenario II. Consider a different model, containing the two chiral representations $(0, \frac{1}{2})$ and $(\frac{1}{2}, 0)$. In the helicity $+1/2$ sector, there are two spin-1/2 nucleons N_A, N_B which are linear combinations of the two chiral representations, with a mixing angle θ

$$\lambda = +\frac{1}{2} : N_A = \cos\theta(0_L, \frac{1}{2}_R) + \sin\theta(\frac{1}{2}_L, 0_R) \quad (48)$$

$$N_B = -\sin\theta(0_L, \frac{1}{2}_R) + \cos\theta(\frac{1}{2}_L, 0_R) \quad (49)$$

We take these states to be mass eigenstates, with eigenvalues m_A, m_B . They are related to the chiral singlet and chiral quartet terms as follows. Consider the squared mass matrix in the basis of the irreducible representations $(\frac{1}{2}, 0)$ and $(0, \frac{1}{2})$

$$m_{\lambda=+1/2}^2 = \begin{pmatrix} \mu_L^2 & \alpha \\ \alpha & \mu_R^2 \end{pmatrix} \quad (50)$$

where $\mu_{L,R}^2$ are the diagonal matrix elements of the chiral singlet $\hat{m}_0^2$, and α is the off-diagonal matrix element of the chiral quartet term in the mass-squared operator of eq. (36). Their relation to the hadronic parameters is

$$\mu_L^2 = m_A^2 \sin^2 \theta + m_B^2 \cos^2 \theta, \mu_R^2 = m_A^2 \cos^2 \theta + m_B^2 \sin^2 \theta, \alpha = \frac{1}{2}(m_A^2 - m_B^2) \sin 2\theta \quad (51)$$

The helicity $-1/2$ sector can be constructed from these states by applying the normality transformation, as explained above. Choosing the two nucleons to have opposite parities $\pi_{N_A} = +1, \pi_{N_B} = -1$, the representation content of the helicity $-1/2$ states is

$$\lambda = -\frac{1}{2} : \quad \Pi N_A = \sin \theta (0_L, \frac{1}{2}_R) + \cos \theta (\frac{1}{2}_L, 0_R) \quad (52)$$

$$\Pi N_B = \cos \theta (0_L, \frac{1}{2}_R) - \sin \theta (\frac{1}{2}_L, 0_R) \quad (53)$$

Next consider the couplings of these states to pions. The pion coupling operator $X^a = \frac{1}{2}(I_R^a - I_L^a)$ is related to the generators of the symmetry, and thus couples only states in the same chiral representations. The simplest way to derive the pion couplings is to consider the transitions $p_{i\uparrow} \rightarrow n_{j\uparrow} \pi^+$, which are mediated by the operator $X^- = X^1 - iX^2$. For example, the diagonal pion transition $N_A \rightarrow N_A \pi$ is computed as

$$\begin{aligned} A(p_{A\uparrow} \rightarrow n_{A\uparrow} \pi^+) &\sim \\ &\left(\cos \theta \langle (1,0)_L, (\frac{1}{2}, -\frac{1}{2})_R | + \sin \theta \langle (\frac{1}{2}, -\frac{1}{2})_L, (1,0)_R | \right) \\ &\times X^- \left(\cos \theta | (1,0)_L, (\frac{1}{2}, +\frac{1}{2})_R \rangle + \sin \theta | (\frac{1}{2}, +\frac{1}{2})_L, (1,0)_R \rangle \right) \\ &= \cos^2 \theta \langle (\frac{1}{2}, -\frac{1}{2})_R | I_R^- | (\frac{1}{2}, +\frac{1}{2})_R \rangle - \sin^2 \theta \langle (\frac{1}{2}, -\frac{1}{2})_L | I_L^- | (\frac{1}{2}, +\frac{1}{2})_L \rangle = \cos 2\theta \end{aligned} \quad (54)$$

where in the last line we show only the chiral isospin (and its third component) changed by the $I_{L,R}^-$ operator. According to the effective Lagrangian of eq. (16), this amplitude is proportional to the coupling $c_2 + c_4$. In a similar way one can obtain all the other couplings in this effective Lagrangian, with the results

$$c_2 + c_4 \sim \cos 2\theta \quad (55)$$

$$c_2 - c_4 \sim -\cos 2\theta \quad (56)$$

$$1 - c_3 \sim -\sin 2\theta \quad (57)$$

This illustrates how a special chiral representation content in the framework of the mended chiral symmetry can lead to specific relations among pion couplings in the effective Lagrangian. In particular, choosing $\theta = 3\pi/4$ reproduces the predictions of the chiral doubling model with linearly transforming baryons ($c_{2-4} = 0$), but without mass degeneracy. The mass parameters of this model are $\mu_L^2 = \mu_R^2 = (m_A^2 + m_B^2)/2$ and $\alpha = -(m_A^2 - m_B^2)/2$, with nondegenerate baryons $m_A^2 - m_B^2 \neq 0$.

Of course, any such explanation leaves open the dynamical origin of this (or any other) particular value of the mixing angle, as well as the representation content. More

complicated representation contents are also possible, e.g. including also $(I_L, I_R) = (1, 1/2), (1/2, 1)$ as in Scenario 1. Such questions could be addressed for example using lattice QCD methods, as discussed recently in Ref. [60].

CONCLUSION AND DISCUSSION

Although parity doubling was observed in the baryon spectrum more than 35 years ago, a complete understanding of its dynamical origin is still lacking. In this paper we both evaluate the empirical evidence for parity doubling and consider some possible theoretical explanations. We propose a new mechanism, based on the dynamical suppression of the $U(1)_A$ violation, and point out a prediction that distinguishes it from $SU(2)_L \times SU(2)_R$ restoration.

Most of the evidence previously offered in favor of parity doubling has been obtained by visually grouping states into multiplets. Given the number of broad, closely spaced states in each channel, this procedure is both ambiguous and imprecise. It does not take into account the resonance widths, nor the fact that different resonances are established with varying degrees of certainty. We propose in this paper a quantitative measure of parity doubling, which is free of these problems. First, we introduce an overlap integral $\Omega_{IJS}(\{C\})$ of the spectral densities in channels with the same spin J , isospin I and strangeness S , but opposite parity. This is computed on an ensemble of control samples (each of them denoted by $\{C\}$), obtained from the real world by arbitrarily permuting the parities of the hadronic states. Finally, the value of the overlap integral Ω_{IJS} in the real world is compared against the statistical distribution on the ensemble, which assigns a probability measure (p -value) to the statement of parity doubling.

The results of our study gives strong evidence for parity doubling in the nonstrange baryon sector (both N and Δ), with spins from $J = 1/2$ to $7/2$. For larger spins the number of states decreases, and we do not expect the control samples to be representative statistically. Still, there is good evidence for parity doubling among high spin, non-strange baryons as can be seen from Table 2. On the other hand, the correlations between positive and negative parity *strange* baryons are not significant. However, some of the states with strangeness expected in the quark model have not yet been seen, which could partly be responsible for the weaker correlations observed in these sectors. A new experimental attack on baryon spectroscopy could shed significant light on the systematics and origins of parity doubling.

In principle our statistical test could be used to search for more complex correlations, such as those expected in models of “chiral symmetry restoration”. In such schemes, additional degeneracies involving hadrons of different isospins (but the same spin) are expected on top of parity doubling. The pattern of the mass spectrum in this case depends on the representation, (j_L, j_R) , of $SU(2)_L \times SU(2)_R$ assumed to be realized. For example, in the $(1/2, 1) \oplus (1, 1/2)$ scheme of Ref. [54] one expects the baryons to be grouped into “quartets” of the form $(N_J^+, N_J^-, \Delta_J^+, \Delta_J^-)$ with the same spin J . We have not used our statistical method to look for more complicated correlations. This would require higher dimensional integrals analogous to eq. (3). In any case no general correlations of this form are observed by eye. However, it is possible that more complicated chiral

representations may not appear uniformly in the spectrum, but be localized in some mass region. This would render our method inapplicable, due to the sparseness of the corresponding control sample.

We have examined a number of possible explanations of parity doubling. First we have tried to clarify the relationship between the usual $SU(2)_L \times SU(2)_R$ chiral symmetry of QCD and parity doubling. The terminology used in the literature discussion has been confusing. In an important sense (for massless u and d quarks) $SU(2)_L \times SU(2)_R$ is not broken at all, but instead represented in the Nambu-Goldstone form by massless pions and isospin multiplets of hadrons that transform non-linearly under chiral transformations. There is nothing to “restore”. Instead, we frame the question as follows: “What is needed in order that baryons fall into Wigner-Weyl representations of $SU(2) \times SU(2)$ with the usual degeneracies and relations among couplings”? The answer, presented in Ref. [11] and reviewed here, is that a set of $SU(2)_L \times SU(2)_R$ *invariant* operator matrix elements must be suppressed, basically amounting to decoupling the pion from the parity doublet states. If this decoupling were complete, it would, in effect, leave *two independent* $SU(2)_L \times SU(2)_R$ symmetries: one under which the pion and non-parity-doubled states transform *non-linearly*, and the other under which the parity doubled states transform *linearly*. A toy model of this type has recently been proposed by Cohen and Glozman [61]. The question is what dynamics are responsible for suppressing these operator matrix elements and giving rise to this additional symmetry? If the dynamics behind effective chiral symmetry restoration applies generically to hadrons high in the spectrum, as advocated, for example, in Ref. [27], then parity doubling should be universal in that domain. It would be puzzling if, for example, some (*e.g.*) highly excited mesons are parity doubled, but some are not. A search for such exceptions would be a particularly effective way to test this explanation of parity doubling.

We emphasize that parity doubling alone is not conclusive evidence for effective chiral symmetry restoration. As discussed in Sec. 3A, chiral symmetry realized linearly on a set of baryons requires that pions decouple from the parity doubled states. In addition, one would have to check the validity of certain relations for axial couplings between the members of the putative chiral multiplet implied by the symmetry algebra implemented in the Wigner-Weyl mode.

Next we considered the possibility that parity doubling is a consequence of dynamical suppression of $U(1)_A$ symmetry breaking in a subsector $\mathfrak{h}$ of the Hilbert space of QCD. Note that this is not equivalent to $U(1)_A$ restoration, as realized for example in large N_c QCD. For example, the effects of the anomaly are still present, and give the η, η' the masses observed in nature. Rather, we require only that the matrix elements of the divergence of the axial singlet current are very small on a subset of baryon states $\langle B_1 | \partial_\mu j_5^\mu | B_2 \rangle \approx 0$, with $B_1, B_2 \in \mathfrak{h}$. This condition is satisfied if the matrix elements of the gluon anomaly term taken between non-strange baryons of opposite parity are very small. The hypothesis that the matrix elements of $\tilde{F}F$ are suppressed among excited baryons may be amenable to study using lattice QCD methods.

Although it predicts parity doublets, the mechanism of dynamical suppression of $U(1)_A$ symmetry breaking does not make predictions for the pion couplings to the doublet states (except that the pions must decouple from parity doubled states at zero momentum, as required by the Goldberger-Treiman relation). Such relations would be obtained, for example, if we required the doublet to be left invariant under $U(1)_A$ and

parity transformations.

In addition to suppression of $U(1)_A$ symmetry violation, we review two other possible origins for parity doubling. First we examine arguments recently put forward that parity doubling in the spectrum can be related to chiral symmetry restoration at short distances, a phenomenon well grounded in perturbative QCD. The only input from short distance chiral symmetry is the superconvergence relation, eq. (26). The rest of the argument requires input from phenomenological models (*e.g.* linear excitation spectra) and large N_c implemented in an unusual way — examining the large n (excitation) limit, but taking $N_c \rightarrow \infty$ first. $\mathcal{O}(1/N_c)$ corrections seem capable of invalidating the conclusions. In any event this exercise provides an example of the extent of additional dynamics required to obtain effective chiral symmetry restoration. Finally we look at the possibility that parity doubling may be a consequence of an intrinsic deformed state that spontaneously violates parity. A mechanism familiar from nuclear and molecular physics, this leads one to expect parity doubling at high spin for both strange and non-strange baryons, but gives little insight into why spin 1/2 and 3/2 nucleons and Δ s should show statistically significant evidence for the symmetry.

An alternative way of obtaining relations among masses and axial couplings which adds dynamical input in a minimal way is offered by the mended chiral symmetry approach of Weinberg [23, 24]. In this approach the set of all pion couplings among baryon states X^a (in sectors of given helicity), together with the isospin operators T^a , are shown to satisfy the $SU(2)_L \times SU(2)_R$ algebra. However, input from Regge theory shows that the mass operator is not a singlet under this symmetry, but contains in addition also a chiral 4-vector term.

This implies that the hadrons transform in *reducible* representations of the $SU(2)_L \times SU(2)_R$ symmetry. Models of hadrons can be constructed by postulating a representation content, and the corresponding mixing angles. Even minimal models containing one or two such representations can reproduce the properties of the lowest lying baryons remarkably well [23, 24].

As examples of this approach we presented two such models. First, to introduce the ideas we describe a model containing a $(N_{1/2}, \Delta_{3/2})$ degenerate pair of the same parity, which reproduces the quark model predictions for pion couplings. This model was previously considered in Refs. [55, 57]. The second model is new and contains two nucleons of opposite parity and spins $J = 1/2$. Their masses and couplings are determined by just three mass parameters and one mixing angle among chiral representations. We show that, for a particular value of the mixing angle, the pion couplings to the parity doublet take precisely the same values as those expected if chiral symmetry were realized linearly on the two states, but without mass degeneracy in the spectrum. This illustrates a natural realization of such relations which does not require chiral symmetry restoration.

ACKNOWLEDGMENTS

We thank W. Bardeen, S. Beane, D. Bugg, T. Cohen, M. Golterman, L. Glozman, A. Manohar, M. Shifman and A. Vainshtein for conversations and correspondence relating to this work. This work is supported in part by funds provided by the U.S. Depart-

ment of Energy (D.O.E.) under cooperative research agreement DE-FC02-94ER40818.

REFERENCES

1. V. Barger and D. Cline, Phys. Lett. **B 26**, 85 (1967); Phys. Rev. Lett. **20** 298 (1968). See also Y. Iwasaki, Prog. Theor. Phys. **44**, 777 (1970), and L. Y. Glozman, arXiv:hep-ph/0105225.
2. See, for example, F. Iachello, Phys. Rev. Lett. **62**, 2440 (1989), D. Robson, Phys. Rev. Lett. **63**, 1890 (1989), and F. Iachello, Phys. Rev. Lett. **63**, 1891 (1989).
3. M. Kirchbach, Mod. Phys. Lett. A **12**, 3177 (1997) [arXiv:nucl-th/9712072].
4. L. Y. Glozman, Phys. Lett. B **475**, 329 (2000) [arXiv:hep-ph/9908207]. L. Y. Glozman, Phys. Lett. B **539**, 257 (2002) [arXiv:hep-ph/0205072].
5. T. D. Cohen and L. Y. Glozman, Phys. Rev. D **65**, 016006 (2002) [arXiv:hep-ph/0102206].
6. T. D. Cohen and L. Y. Glozman, Int. J. Mod. Phys. A **17**, 1327 (2002) [arXiv:hep-ph/0201242]. L. Y. Glozman, Acta Phys. Polon. B **35**, 2985 (2004) [arXiv:hep-ph/0410194].
7. A. E. Inopin, arXiv:hep-ph/0404048.
8. M. Shifman, arXiv:hep-ph/0507246.
9. D. Jido, T. Hatsuda and T. Kunihiro, Phys. Rev. Lett. **84**, 3252 (2000) [arXiv:hep-ph/9910375].
10. For a summary with further references, see D. Jido, M. Oka and A. Hosaka, Prog. Theor. Phys. **106**, 873 (2001) [arXiv:hep-ph/0110005].
11. R. L. Jaffe, D. Pirjol and A. Scardicchio, arXiv:hep-ph/0511081(v2), to be published in Phys. Rev. Lett.
12. S. Weinberg, Physica, **96A**, 327 (1979); J. Gasser and H. Leutwyler, Nucl. Phys. B **250**, 465 (1985).
13. S. Weinberg, Phys. Rev. **166**, 1568 (1968).
14. S. Weinberg, *The Quantum Theory of Fields, Vol. 2: Modern Applications*.
15. S. R. Coleman, J. Wess and B. Zumino, Phys. Rev. **177**, 2239 (1969); C. G. Callan, S. R. Coleman, J. Wess and B. Zumino, Phys. Rev. **177**, 2247 (1969).
16. C. DeTar and T. Kunihiro, Phys. Rev. D **39**, 2805 (1989).
17. For a review, see G. 't Hooft, Phys. Rept. **142**, 357 (1986).
18. S. Coleman, J. Math. Phys. **7**, 787 (1966).
19. D. J. Broadhurst, Phys. Lett. B **101**, 423 (1981).
20. To our knowledge, the possibility that parity doubling in QCD is a consequence of the suppression of $U(1)_A$ violating matrix elements between baryons was first suggested in [21].
21. R. L. Jaffe, at the Workshop on Key Issues in Hadronic Physics, Duck, North Carolina, November 2000.
22. K. Johnson and C. B. Thorn, Phys. Rev. D **13**, 1934 (1976).
23. S. Weinberg, Phys. Rev. **177**, 2604 (1969).
24. S. Weinberg, Phys. Rev. Lett. **65**, 1177 (1990).
25. R. F. Dashen and A. V. Manohar, Phys. Lett. B **315**, 425 (1993) [arXiv:hep-ph/9307241]; Phys. Lett. B **315**, 438 (1993) [arXiv:hep-ph/9307242].
26. R. F. Dashen, E. Jenkins and A. V. Manohar, Phys. Rev. D **49**, 4713 (1994) [Erratum-ibid. D **51**, 2489 (1995)] [arXiv:hep-ph/9310379]; Phys. Rev. D **51**, 3697 (1995) [arXiv:hep-ph/9411234].
27. L. Y. Glozman, Int. J. Mod. Phys. A **21**, 475 (2006) [arXiv:hep-ph/0411281].
28. E. Klempt, Phys. Lett. B **559**, 144 (2003) [arXiv:hep-ph/0212241].
29. S. Eidelman *et al.* [Particle Data Group], Phys. Lett. B **592** (2004) 1.
30. See Ref. [29]. The table of interest can be found at http://pdg.lbl.gov/2005/reviews/ndelmini_b861.pdf
31. D. V. Bugg, Phys. Rept. **397**, 257 (2004) [arXiv:hep-ex/0412045].
32. R. F. Dashen, Phys. Rev. **183**, 1245 (1969).
33. I. I. Kogan, A. Kovner and M. A. Shifman, Phys. Rev. D **59**, 016001 (1999) [arXiv:hep-ph/9807286].
34. M. Knecht and J. Stern, arXiv:hep-ph/9411253. J. Stern, arXiv:hep-ph/9712438.
35. D. Weingarten, Phys. Rev. Lett. **51** 1830 (1983); E. Witten, Phys. Rev. Lett. **51**, 2351 (1983); S. Nussinov, Phys. Rev. Lett. **52**, 966 (1984); D. Espriu, M. Gross and J. F. Wheeler, Phys. Lett. **146 B**, 67 (1984).
36. G. M. Shore and G. Veneziano, Phys. Lett. B **244**, 75 (1990); G. M. Shore and G. Veneziano, Nucl.

- Phys. B **381**, 23 (1992). For a recent update of the experimental data see G. M. Shore, arXiv:hep-ph/0601051.
37. N. Kochelev and D. P. Min, arXiv:hep-ph/0508288.
 38. D. Diakonov, M. V. Polyakov and C. Weiss, Nucl. Phys. B **461**, 539 (1996) [arXiv:hep-ph/9510232].
 39. W. A. Bardeen and C. T. Hill, Phys. Rev. D **49**, 409 (1994) [arXiv:hep-ph/9304265]. W. A. Bardeen, E. J. Eichten and C. T. Hill, Phys. Rev. D **68**, 054024 (2003) [arXiv:hep-ph/0305049].
 40. S. Weinberg, Phys. Rev. Lett. **18**, 507 (1967).
 41. G. 't Hooft, Phys. Rev. Lett. **37**, 8 (1976).
 42. J. Engel, M. Bender, J. Dobaczewski, J. H. De Jesus and P. Olbratowski, Phys. Rev. C **68**, 025501 (2003) [arXiv:nucl-th/0304075].
 43. For a review of reflection asymmetric nuclei, see P. A. Butler, Act. Phys. Pol. **B29**, 289 (1998).
 44. J. F. C. Cocks *et al.*, Phys. Rev. Lett. **78** 2820 (1997).
 45. For a recent re-examination of this idea, see Ref. [46]
 46. F. Wilczek, arXiv:hep-ph/0409168, and A. Selem and F. Wilczek, in preparation.
 47. S. R. Beane, Phys. Rev. D **64**, 116010 (2001) [arXiv:hep-ph/0106022].
 48. M. Shifman, private correspondence.
 49. M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **147**, 385 (1979).
 50. M. Golterman and S. Peris, Phys. Rev. D **67** 096001 (2003) [arXiv:hep-ph/0207060]; O. Cata, M. Golterman, and S. Peris [arXiv:hep-ph/0602194].
 51. G. 't Hooft, Nucl. Phys. B **75**, 461 (1974).
 52. B. Blok, M. A. Shifman and D. X. Zhang, Phys. Rev. D **57**, 2691 (1998) [Erratum-ibid. D **59**, 019901 (1999)] [arXiv:hep-ph/9709333].
 53. S. L. Adler, Phys. Rev. Lett. **14**, 1051 (1965); W. I. Weisberger, Phys. Rev. Lett. **14**, 1047 (1965).
 54. D. Jido, T. Hatsuda and T. Kunihiro, Phys. Rev. Lett. **84**, 3252 (2000) [arXiv:hep-ph/9910375].
 55. S. Weinberg, arXiv:hep-ph/9412326.
 56. S. R. Beane, arXiv:hep-ph/9512228;
 57. S. R. Beane, Phys. Rev. D **59**, 031901 (1999) [arXiv:hep-ph/9809328];
 58. S. R. Beane and M. J. Savage, Phys. Lett. B **556**, 142 (2003) [arXiv:hep-ph/0212106];
 59. S. R. Beane, Phys. Rev. D **70**, 114010 (2004) [arXiv:hep-ph/0408066].
 60. S. R. Beane and M. J. Savage, Phys. Rev. D **70**, 074029 (2004) [arXiv:hep-ph/0404131].
 61. T. Cohen and L. Glozman, hep-ph/0512185.